

Rabbit anti-Human, Mouse XPO1 Polyclonal Antibody | anti-XPO1 antibody

XPO1 Antibody

CATALOG # AAA232728	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	-----------------------------------	--


PRODUCT SYNONYMS

XPO1; Polyclonal Antibody; XPO1 Antibody; Exportin-1; Exp1; Chromosome region maintenance 1 protein homolog; CRM1; anti-XPO1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	1071
Applicable Applications for anti-XPO1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Exportin-1 protein (625-845AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-5595 / sc-7825 / sc-7826
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of HeLa cells using AAA232728 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)




IHC (Immunohistochemistry)

IHC image of AAA232728 diluted at 1:150 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA232728 diluted at 1:150 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: K562 whole cell lysate,NIH/3T3 whole cell lysateAll lanes: XPO1 antibody at 3ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 124 KDaObserved band size: 124 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-XPO1 antibody

Mediates the nuclear export of cellular proteins (cargos) bearing a leucine-rich nuclear export signal (NES) and of RNAs. In the nucleus, in association with RANBP3, binds cooperatively to the NES on its target protein and to the GTPase RAN in its active GTP-bound form (Ran-GTP). Docking of this complex to the nuclear pore complex (NPC) is mediated through binding to nucleoporins. Upon transit of a nuclear export complex into the cytoplasm, disassembling of the complex and hydrolysis of Ran-GTP to Ran-GDP (induced by RANBP1 and RANGAP1, respectively) cause release of the cargo from the export receptor. The directionality of nuclear export is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. Involved in U3 snoRNA transport from Cajal bodies to nucleoli. Binds to late precursor U3 snoRNA bearing a TMG cap. Several viruses, among them HIV-1, HTLV-1 and influenza A use it to export their unspliced or incompletely spliced RNAs out of the nucleus. Interacts with, and mediates the nuclear export of HIV-1 Rev and HTLV-1 Rex proteins. Involved in HTLV-1 Rex multimerization.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507943
NCBI GeneID	7514
NCBI Accession #	NP_003391.1
NCBI GenBank Nucleotide #	NM_003400.3
UniProt Accession #	O14980; Q63HP8; Q68CP3; Q99433; A6NL14; A8K1K5; D6W5E2
Molecular Weight	123,386 Da
NCBI Official Full Name	exportin-1
NCBI Official Synonym Full Names	exportin 1
NCBI Official Symbol	XPO1
NCBI Official Synonym Symbols	emb; CRM1; exp1
NCBI Protein Information	exportin-1
UniProt Protein Name	Exportin-1
UniProt Gene Name	XPO1
UniProt Synonym Gene Names	CRM1; Exp1
UniProt Entry Name	XPO1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/xpo1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/232728>