

Rabbit anti-Human, Mouse DDR2 Polyclonal Antibody | anti-DDR2 antibody

DDR2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232870	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

DDR2; Polyclonal Antibody; DDR2 Antibody; Discoidin domain-containing receptor 2; Discoidin domain receptor 2; CD167 antigen-like family member B; Discoidin domain-containing receptor tyrosine kinase 2; Neurotrophic tyrosine kinase; receptor-related 3; Receptor protein-tyrosine kinase TKT; Tyrosine-protein kinase TYRO10; CD167b; NTRKR3; TKT; TYRO10; anti-DDR2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	855
Applicable Applications for anti-DDR2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Discoidin domain-containing receptor 2 protein (447-540AA)
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Conjugate	Non-conjugated
Species	Human
Research Area	Neuroscience
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-7555 / sc-8989
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES



IHC (Immunohistochemisry)
Immunofluorescent analysis of HepG2 cells using AAA232870 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded human heart tissue using AAA232870 at dilution of 1:100



WB (Western Blot)
Western BlotPositive WB detected in: HeLa whole cell lysate,Jurkat whole cell lysate,HEK293 whole cell lysate,NIH/3T3 whole cell lysate,A549 whole cell lysateAll lanes: DDR2 antibody at 3.5ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 97 KDaObserved band size: 97 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-DDR2 antibody	Tyrosine kinase that functions as cell surface receptor for fibrillar collagen and regulates cell differentiation, remodeling of the extracellular matrix, cell migration and cell proliferation. Required for normal bone development. Regulates osteoblast differentiation and chondrocyte maturation via a signaling pathway that involves MAP kinases and leads to the activation of the transcription factor RUNX2. Regulates remodeling of the extracellular matrix by up-regulation of the collagenases MMP1, MMP2 and MMP13, and thereby facilitates cell migration and tumor cell invasion. Promotes fibroblast migration and proliferation, and thereby contributes to cutaneous wound healing.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	62420886
NCBI GeneID	4921
NCBI Accession #	NP_001014796.1
NCBI GenBank Nucleotide #	NM_001014796.1
UniProt Accession #	Q16832; Q7Z730
NCBI Official Full Name	discoidin domain-containing receptor 2
NCBI Official Synonym Full Names	discoidin domain receptor tyrosine kinase 2
NCBI Official Symbol	DDR2

NCBI Official Synonym Symbols	TKT; MIG20a; NTRKR3; TYRO10
NCBI Protein Information	discoidin domain-containing receptor 2
UniProt Protein Name	Discoidin domain-containing receptor 2
UniProt Gene Name	DDR2
UniProt Synonym Gene Names	NTRKR3; TKT; TYRO10; Discoidin domain receptor 2
UniProt Entry Name	DDR2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ddr2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/232870>