

Rabbit anti-Human MYO7A Polyclonal Antibody | anti-MYO7A antibody

MYO7A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232873	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

MYO7A; Polyclonal Antibody; MYO7A Antibody; Unconventional myosin-VIIa; USH1B; anti-MYO7A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	2215
Applicable Applications for anti-MYO7A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Unconventional myosin-VIIa protein (838-968AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-25834 / sc-26709
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES


IHC (Immunohistochemisry)

Immunofluorescent analysis of HepG2 cells using AAA232873 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)


IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded human adrenal gland tissue using AAA232873 at dilution of 1:100



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human liver cancer using AAA232873 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-MYO7A antibody	Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Their highly divergent tails bind to membranous compartments, which are then moved relative to actin filaments. In the retina, plays an important role in the renewal of the outer photoreceptor disks. Plays an important role in the distribution and migration of retinal pigment epithelial (RPE) melanosomes and phagosomes, and in the regulation of opsin transport in retinal photoreceptors. In the inner ear, plays an important role in differentiation, morphogenesis and organization of cochlear hair cell bundles. Involved in hair-cell vesicle trafficking of aminoglycosides, which are known to induce ototoxicity (By similarity). Motor protein that is a part of the functional network formed by USH1C, USH1G, CDH23 and MYO7A that mediates mechanotransduction in cochlear hair cells. Required for normal hearing.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	189083798
NCBI GeneID	4647
NCBI Accession #	NP_000251.3
NCBI GenBank Nucleotide #	NM_000260.3
UniProt Accession #	Q13402; P78427; Q13321; Q14785; Q92821; Q92822; B9A011; F8VUN5
Molecular Weight	249,165 Da
NCBI Official Full Name	unconventional myosin-VIIa isoform 1
NCBI Official Synonym Full Names	myosin VIIA
NCBI Official Symbol	MYO7A
NCBI Official Synonym Symbols	DFNB2; MYU7A; NSRD2; USH1B; DFNA11; MYOVIIA
NCBI Protein Information	unconventional myosin-VIIa
UniProt Protein Name	Unconventional myosin-VIIa
UniProt Gene Name	MYO7A
UniProt Synonym Gene Names	USH1B
UniProt Entry Name	MYO7A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.