

Rabbit anti-Human RACGAP1 Polyclonal Antibody | anti-RACGAP1 antibody

RACGAP1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA233092	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

RACGAP1; Polyclonal Antibody; RACGAP1 Antibody; Rac GTPase-activating protein 1; Male germ cell RacGap; MgcRacGAP; Protein CYK4 homolog; CYK4; HsCYK-4; anti-RACGAP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	632
Applicable Applications for anti-RACGAP1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Rac GTPase-activating protein 1 protein (86-331AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-82733 / sc-82735 / sc-98617
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of HepG2 cells using AAA233092 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human thymus tissue using AAA233092 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-RACGAP1 antibody	Component of the centralspindlin complex that serves as a microtubule-dependent and Rho-mediated signaling required for the myosin contractile ring formation during the cell cycle cytokinesis. Required for proper attachment of the midbody to the cell membrane during cytokinesis. Plays key roles in controlling cell growth and differentiation of hematopoietic cells through mechanisms other than regulating Rac GTPase activity. Also involved in the regulation of growth-related processes in adipocytes and myoblasts. May be involved in regulating spermatogenesis and in the RACGAP1 pathway in neuronal proliferation. Shows strong GAP (GTPase activation) activity towards CDC42 and RAC1 and less towards RHOA. Essential for the early stages of embryogenesis. May play a role in regulating cortical activity through RHOA during cytokinesis. May participate in the regulation of sulfate transport in male germ cells.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	186910300
NCBI GeneID	29127
NCBI Accession #	NP_001119575.1
NCBI GenBank Nucleotide #	NM_001126103.2
UniProt Accession #	Q9H0H5; Q6PJ26; Q9NWN2; Q9P250; Q9P2W2
Molecular Weight	71,027 Da
NCBI Official Full Name	rac GTPase-activating protein 1 isoform a
NCBI Official Synonym Full Names	Rac GTPase activating protein 1
NCBI Official Symbol	RACGAP1
NCBI Official Synonym Symbols	CYK4; ID-GAP; HsCYK-4; MgcRacGAP
NCBI Protein Information	rac GTPase-activating protein 1
UniProt Protein Name	Rac GTPase-activating protein 1
UniProt Gene Name	RACGAP1
UniProt Synonym Gene Names	MgcRacGAP; CYK4; HsCYK-4
UniProt Entry Name	RGAP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/racgap1-antibody-immunofluorescence-immunohistochemistry-elisa/233092>