

Rabbit anti-Human SIRT7 Polyclonal Antibody | anti-SIRT7 antibody

SIRT7 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA233098	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

SIRT7; Polyclonal Antibody; SIRT7 Antibody; NAD-dependent protein deacetylase sirtuin-7; Regulatory protein SIR2 homolog 7; SIR2-like protein 7; SIR2L7; anti-SIRT7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	400
Applicable Applications for anti-SIRT7 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human NAD-dependent protein deacetylase sirtuin-7 protein (242-379AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-66281 / sc-66283
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of Hela cells using AAA233098 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney tissue using AAA233098 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-SIRT7 antibody	NAD-dependent protein deacetylase that specifically mediates deacetylation of histone H3 at 'Lys-18' (H3K18Ac). In contrast to other histone deacetylases, displays selectivity for a single histone mark, H3K18Ac, directly linked to control of gene expression. H3K18Ac is mainly present around the transcription start site of genes and has been linked to activation of nuclear hormone receptors. SIRT7 thereby acts as a transcription repressor. Moreover, H3K18 hypoacetylation has been reported as a marker of malignancy in various cancers and seems to maintain the transformed phenotype of cancer cells. These data suggest that SIRT7 may play a key role in oncogenic transformation by suppresses expression of tumor suppressor genes by locus-specific deacetylation of H3K18Ac at promoter regions. Also required to restore the transcription of ribosomal RNA (rRNA) at the exit from mitosis: promotes the association of RNA polymerase I with the rDNA promoter region and coding region. Stimulates transcription activity of the RNA polymerase I complex. May also deacetylate p53/TP53 and promotes cell survival, however such data need additional confirmation.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	7706712
NCBI GeneID	51547
NCBI Accession #	NP_057622.1
NCBI GenBank Nucleotide #	NM_016538.2
UniProt Accession #	Q9NRC8; Q3MIK4; Q9NSZ6; Q9NUS6; A8K2K0; B3KSU8
Molecular Weight	35,851 Da
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-7
NCBI Official Synonym Full Names	sirtuin 7
NCBI Official Symbol	SIRT7
NCBI Official Synonym Symbols	SIR2L7
NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-7
UniProt Protein Name	NAD-dependent protein deacetylase sirtuin-7
UniProt Gene Name	SIRT7
UniProt Synonym Gene Names	SIR2L7
UniProt Entry Name	SIR7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/sirt7-antibody-immunofluorescence-immunohistochemistry-elisa/233098>