

Rabbit anti-Human TMEM173 Polyclonal Antibody | anti-TMEM173 antibody

TMEM173 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA233301	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

TMEM173; Polyclonal Antibody; TMEM173 Antibody; Stimulator of interferon genes protein1; Endoplasmic reticulum interferon stimulator; ERIS; Mediator of IRF3 activation; MITA; anti-TMEM173 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	379
Applicable Applications for anti-TMEM173 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Stimulator of interferon genes protein (207-379AA)
Conjugation	Non-conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemisry)

Immunofluorescent analysis of MCF-7 cells using AAA233301 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohiostchemistry)

IHC image of AAA233301 diluted at 1:400 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: MCF-7 whole cell lysate All lanes: TMEM173 antibody at 2.7ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 43 KDa Observed band size: 43 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-TMEM173 antibody

Facilitator of innate immune signaling that acts as a sensor of cytosolic DNA from bacteria and viruses and promotes the production of type I interferon (IFN-alpha and IFN-beta). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm. Acts by recognizing and binding cyclic di-GMP (c-di-GMP), a second messenger produced by bacteria, and cyclic GMP-AMP (cGAMP), a messenger produced in response to DNA virus in the cytosol: upon binding of c-di-GMP or cGAMP, autoinhibition is alleviated and TMEM173/STING is able to activate both NF-kappa-B and IRF3 transcription pathways to induce expression of type I interferon and exert a potent anti-viral state. May be involved in translocon function, the translocon possibly being able to influence the induction of type I interferons. May be involved in transduction of apoptotic signals via its association with the major histocompatibility complex class II (MHC-II). Mediates death signaling via activation of the extracellular signal-regulated kinase (ERK) pathway. Essential for the induction of IFN-beta in response to human herpes simplex virus 1 (HHV-1) infection. Exhibits 2', 3' phosphodiester linkage-specific ligand recognition. Can bind both 2'-3' linked cGAMP and 3'-3' linked cGAMP but is preferentially activated by 2'-3' linked cGAMP (PubMed:26300263).

NCBI AND UNIPROT INFORMATION

NCBI GI #	38093659
NCBI GeneID	340061
NCBI Accession #	NP_938023.1
NCBI GenBank Nucleotide #	NM_198282.3
UniProt Accession #	Q86WV6; A8K3P6; B6EB35; D6RBX0; D6RE01; D6RID9
Molecular Weight	42,193 Da
NCBI Official Full Name	stimulator of interferon genes protein isoform 1
NCBI Official Synonym Full Names	transmembrane protein 173
NCBI Official Symbol	TMEM173
NCBI Official Synonym Symbols	ERIS; MITA; MPYS; SAVI; NET23; STING; hMITA; hSTING
NCBI Protein Information	stimulator of interferon genes protein
UniProt Protein Name	Stimulator of interferon genes protein
UniProt Gene Name	TMEM173
UniProt Synonym Gene Names	; hMITA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tmem173-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/233301>