

Rabbit anti-Human MST1 Polyclonal Antibody | anti-MST1 antibody

MST1 Antibody

CATALOG # AAA234087	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

MST1; Polyclonal Antibody; MST1 Antibody; Hepatocyte growth factor-like protein; Macrophage stimulatory protein; Macrophage-stimulating protein; MSP; Hepatocyte growth factor-like protein alpha chain; Hepatocyte growth factor-like protein beta chain; D3F15S2; DNF15S2; HGFL; anti-MST1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	725
Applicable Applications for anti-MST1 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Hepatocyte growth factor-like protein (363-465AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of AAA234087 diluted at 1:400 and staining in paraffin-embedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234087 diluted at 1:400 and staining in paraffin-embedded human gastric cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-MST1 antibody	extracellular exosome, extracellular region, extracellular space, receptor tyrosine kinase binding, hepatocyte growth factor receptor signaling pathway, negative regulation of gluconeogenesis, regulation of cAMP-dependent protein kinase activity
Product Categories/Family for anti-MST1 antibody	Cancer; Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	205277383
NCBI GeneID	4485
NCBI Accession #	NP_066278.3
NCBI GenBank Nucleotide #	NM_020998.3
UniProt Accession #	P26927; Q13350; Q14870; Q6GTN4; A6NLA3; A8MSX3
Molecular Weight	80,320 Da
NCBI Official Full Name	hepatocyte growth factor-like protein
NCBI Official Synonym Full Names	macrophage stimulating 1
NCBI Official Symbol	MST1
NCBI Official Synonym Symbols	MSP; HGFL; NF15S2; D3F15S2; DNF15S2
NCBI Protein Information	hepatocyte growth factor-like protein
UniProt Protein Name	Hepatocyte growth factor-like protein
UniProt Gene Name	MST1
UniProt Synonym Gene Names	D3F15S2; DNF15S2; HGFL; MSP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mst1-antibody-immunohistochemistry-elisa/234087>