

Rabbit anti-Human PCMT1 Polyclonal Antibody | anti-PCMT1 antibody

PCMT1 Antibody

CATALOG # AAA234096	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

PCMT1; Polyclonal Antibody; PCMT1 Antibody; Protein-L-isoaspartate(D-aspartate) O-methyltransferase; PIMT; L-isoaspartyl protein carboxyl methyltransferase; Protein L-isoaspartyl/D-aspartyl methyltransferase; Protein-beta-aspartate methyltransferase; anti-PCMT1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	286
Applicable Applications for anti-PCMT1 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Protein-L-isoaspartate(D-aspartate) O-methyltransferase protein (117-228AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of AAA234096 diluted at 1:200 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234096 diluted at 1:200 and staining in paraffin-embedded human pancreatic tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-PCMT1 antibody	Catalyzes the methyl esterification of L-isoaspartyl and D-aspartyl residues in peptides and proteins that result from spontaneous decomposition of normal L-aspartyl and L-asparaginyl residues. It plays a role in the repair and/or degradation of damaged proteins. Acts on EIF4EBP2, microtubule-associated protein 2, calreticulin, clathrin light chains a and b, Ubiquitin carboxyl-terminal hydrolase isozyme L1, phosphatidylethanolamine-binding protein 1, stathmin, beta-synuclein and alpha-synuclein.
Product Categories/Family for anti-PCMT1 antibody	Cell Biology

NCBI AND UNIPROT INFORMATION

NCBI GI #	354983493
NCBI GeneID	5110
NCBI Accession #	NP_001238978.1
NCBI GenBank Nucleotide #	NM_001252049.1
UniProt Accession #	P22061; Q14661; Q16556; Q5VYC1; Q5VYC2; Q93061; Q96II9; Q99625; Q9BQV7; Q9BQV8; A8K109; J3KP72
Molecular Weight	24,679 Da
NCBI Official Full Name	protein-L-isoaspartate(D-aspartate) O-methyltransferase isoform 2
NCBI Official Synonym Full Names	protein-L-isoaspartate (D-aspartate) O-methyltransferase
NCBI Official Symbol	PCMT1
NCBI Official Synonym Symbols	PIMT
NCBI Protein Information	protein-L-isoaspartate(D-aspartate) O-methyltransferase
UniProt Protein Name	Protein-L-isoaspartate(D-aspartate) O-methyltransferase
UniProt Gene Name	PCMT1
UniProt Synonym Gene Names	PIMT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pcmt1-antibody-immunohistochemistry-elisa/234096>