

Rabbit anti-Human TCP1 Polyclonal Antibody | anti-TCP1 antibody

TCP1 Antibody

CATALOG # AAA234115	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TCP1; Polyclonal Antibody; TCP1 Antibody; T-complex protein 1 subunit alpha; TCP-1-alpha; CCT-alpha; CCT1; CCTA; anti-TCP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	556
Applicable Applications for anti-TCP1 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human T-complex protein 1 subunit alpha protein (406-506AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of AAA234115 diluted at 1:800 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234115 diluted at 1:800 and staining in paraffin-embedded human liver cancer tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-TCP1 antibody	Molecular chaperone; assists the folding of proteins upon ATP hydrolysis. As part of the BBS/CCT complex may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia. Known to play a role, in vitro, in the folding of actin and tubulin.
Product Categories/Family for anti-TCP1 antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	57863257
NCBI GeneID	6950
NCBI Accession #	NP_110379.2
NCBI GenBank Nucleotide #	NM_030752.2
UniProt Accession #	P17987; Q15556; Q5TCM3; E1P5B2
Molecular Weight	60,344 Da
NCBI Official Full Name	T-complex protein 1 subunit alpha isoform a
NCBI Official Synonym Full Names	t-complex 1
NCBI Official Symbol	TCP1
NCBI Official Synonym Symbols	CCT1; CCTa; D6S230E; CCT-alpha; TCP-1-alpha
NCBI Protein Information	T-complex protein 1 subunit alpha
UniProt Protein Name	T-complex protein 1 subunit alpha
UniProt Gene Name	TCP1
UniProt Synonym Gene Names	CCT1; CCTA; TCP-1-alpha

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tcp1-antibody-immunohistochemistry-elisa/234115>