

Rabbit anti-Human LPP Polyclonal Antibody | anti-LPP antibody

LPP Antibody

CATALOG # AAA234166	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

LPP; Polyclonal Antibody; LPP Antibody; Lipoma-preferred partner; LIM domain-containing preferred translocation partner in lipoma; anti-LPP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	612
Applicable Applications for anti-LPP antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Lipoma-preferred partner protein (227-391AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image of AAA234166 diluted at 1:300 and staining in paraffin-embedded human pancreatic cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234166 diluted at 1:300 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: 293T whole cell lysateAll lanes: LPP antibody at 7.5ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 66 KDaObserved band size: 66 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-LPP antibody	May play a structural role at sites of cell adhesion in maintaining cell shape and motility. In addition to these structural functions, it may also be implicated in signaling events and activation of gene transcription. May be involved in signal transduction from cell adhesion sites to the nucleus allowing successful integration of signals arising from soluble factors and cell-cell adhesion sites. Also suggested to serve as a scaffold protein upon which distinct protein complexes are assembled in the cytoplasm and in the nucleus.
Product Categories/Family for anti-LPP antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	264681524
NCBI GeneID	4026
NCBI Accession #	NP_001161143.1
NCBI GenBank Nucleotide #	NM_001167671.2
UniProt Accession #	Q93052; Q8NFX5; A1L4L6; D3DNV6
Molecular Weight	65,746 Da
NCBI Official Full Name	lipoma-preferred partner isoform a
NCBI Official Synonym Full Names	LIM domain containing preferred translocation partner in lipoma
NCBI Official Symbol	LPP
NCBI Protein Information	lipoma-preferred partner
UniProt Protein Name	Lipoma-preferred partner
UniProt Gene Name	LPP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/lpp-antibody-immunohistochemistry-western-blot-elisa/234166>