

Rabbit anti-Human WASF3 Polyclonal Antibody | anti-WASF3 antibody

WASF3 Antibody

CATALOG # AAA234189	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

WASF3; Polyclonal Antibody; WASF3 Antibody; Wiskott-Aldrich syndrome protein family member 3; WASP family protein member 3; Protein WAVE-3; Verprolin homology domain-containing protein 3; KIAA0900; SCAR3; WAVE3; anti-WASF3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	499
Applicable Applications for anti-WASF3 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Wiskott-Aldrich syndrome protein family member 3 protein (368-502AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of AAA234189 diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234189 diluted at 1:100 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: MCF-7 whole cell lysateAll lanes: WASF3 antibody at 7ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 56 KDaObserved band size: 56 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-WASF3 antibody	Downstream effector molecules involved in the transmission of signals from tyrosine kinase receptors and small GTPases to the actin cytoskeleton. Plays a role in the regulation of cell morphology and cytoskeletal organization. Required in the control of cell shape.
Product Categories/Family for anti-WASF3 antibody	Cancer; Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	634743292
NCBI GeneID	10810
NCBI Accession #	NP_001278894.1
NCBI GenBank Nucleotide #	NM_001291965.1
UniProt Accession #	Q9UPY6; O94974; Q86VQ2
Molecular Weight	55,007 Da
NCBI Official Full Name	wiskott-Aldrich syndrome protein family member 3 isoform 2
NCBI Official Synonym Full Names	WAS protein family member 3
NCBI Official Symbol	WASF3
NCBI Official Synonym Symbols	SCAR3; WAVE3; Brush-1
NCBI Protein Information	wiskott-Aldrich syndrome protein family member 3
UniProt Protein Name	Wiskott-Aldrich syndrome protein family member 3
UniProt Gene Name	WASF3
UniProt Synonym Gene Names	KIAA0900; SCAR3; WAVE3; WASP family protein member 3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/wasf3-antibody-immunohistochemistry-western-blot-elisa/234189>