

Rabbit anti-Human, Rat TNKS Polyclonal Antibody | anti-TNKS antibody

TNKS Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234215	Rabbit	Human, Rat	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

TNKS; Polyclonal Antibody; TNKS Antibody; Tankyrase-1; TANK1; ADP-ribosyltransferase diphtheria toxin-like 5; ARTD5; Poly [ADP-ribose] polymerase 5A; TNKS-1; TRF1-interacting ankyrin-related ADP-ribose polymerase; Tankyrase I; PARP5A; PARPL; TIN1; TINF1; TNKS1; anti-TNKS antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	1327
Applicable Applications for anti-TNKS antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Tankyrase-1 protein (35-130AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

Immunofluorescence staining of HepG2 cells with AAA234215 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of AAA234215 diluted at 1:400 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: Jurkat whole cell lysateAll lanes: TNKS antibody at 4.8ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 143, 68 KDaObserved band size: 143, 68 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-TNKS antibody	Poly-ADP-ribosyltransferase involved in various processes such as Wnt signaling pathway, telomere length and vesicle trafficking. Acts as an activator of the Wnt signaling pathway by mediating poly-ADP-ribosylation (PARsylation) of AXIN1 and AXIN2, 2 key components of the beta-catenin destruction complex: poly-ADP-ribosylated target proteins are recognized by RNF146, which mediates their ubiquitination and subsequent degradation. Also mediates PARsylation of BLZF1 and CASC3, followed by recruitment of RNF146 and subsequent ubiquitination. Mediates PARsylation of TERF1, thereby contributing to the regulation of telomere length. Involved in centrosome maturation during prometaphase by mediating PARsylation of HEPACAM2/MIK1. May also regulate vesicle trafficking and modulate the subcellular distribution of SLC2A4/GLUT4-vesicles. May be involved in spindle pole assembly through PARsylation of NUMA1. Stimulates 26S proteasome activity.
Product Categories/Family for anti-TNKS antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	87239981
NCBI GeneID	8658
NCBI Accession #	NP_003738.2
NCBI GenBank Nucleotide #	NM_003747.2
UniProt Accession #	O95271; O95272; Q4G0F2
Molecular Weight	67,203 Da
NCBI Official Full Name	tankyrase-1
NCBI Official Synonym Full Names	tankyrase
NCBI Official Symbol	TNKS
NCBI Official Synonym Symbols	TIN1; ARTD5; PARPL; TINF1; TNKS1; pART5; PARP5A; PARP-5a
NCBI Protein Information	tankyrase-1
UniProt Protein Name	Tankyrase-1
UniProt Gene Name	TNKS
UniProt Synonym Gene Names	PARP5A; PARPL; TIN1; TINF1; TNKS1; TANK1; ARTD5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.