

Rabbit anti-Human, Mouse H2AFZ Polyclonal Antibody | anti-H2AFZ antibody

H2AFZ (Ab-4) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234360	Rabbit	Human, Mouse	Chromatin Immunoprecipitation, Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

H2AFZ; Polyclonal Antibody; H2AFZ (Ab-4) Antibody; Histone H2A.Z; H2A/z; H2AZ; anti-H2AFZ antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	128
Applicable Applications for anti-H2AFZ antibody	ChIP (Chromatin immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Peptide sequence around site of Lys (4) derived from human Histone H2A.Z.
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells with AAA234360 at 1:10, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).




IHC (Immunohistochemistry)

IHC image of AAA234360 diluted at 1:20 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234360 diluted at 1:20 and staining in paraffin-embedded human prostate tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: K562 whole cell lysate, Mouse spleen tissueAll lanes: H2AFZ antibody at 1ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 14 KDaObserved band size: 14 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-H2AFZ antibody	Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in the formation of constitutive heterochromatin. May be required for chromosome segregation during cell division.
Product Categories/Family for anti-H2AFZ antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504255
NCBI GeneID	3015
NCBI Accession #	NP_002097.1
NCBI GenBank Nucleotide #	NM_002106.3
UniProt Accession #	P0C0S5; P17317; Q6I9U0; B2RD56
Molecular Weight	13,553 Da
NCBI Official Full Name	histone H2A.Z
NCBI Official Synonym Full Names	H2A histone family member Z
NCBI Official Symbol	H2AFZ
NCBI Official Synonym Symbols	H2AZ; H2A.z; H2A/z; H2A.Z-1
NCBI Protein Information	histone H2A.Z
UniProt Protein Name	Histone H2A.Z
UniProt Gene Name	H2AFZ
UniProt Synonym Gene Names	H2AZ; H2A/z

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/h2afz-antibody-chromatin-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/234360>