

Rabbit anti-Human, Rat OLFM4 Polyclonal Antibody | anti-OLFM4 antibody

OLFM4 Antibody

CATALOG # AAA234471	HOST Rabbit	REACTIVITY Human, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	---------------------------------	--


PRODUCT SYNONYMS

OLFM4; Polyclonal Antibody; OLFM4 Antibody; Olfactomedin-4; OLM4; Antiapoptotic protein GW112; G-CSF-stimulated clone 1 protein; hGC-1; hOLfD; GW112; UNQ362/PRO698; anti-OLFM4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	510
Applicable Applications for anti-OLFM4 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Olfactomedin-4 protein (224-334AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image of AAA234471 diluted at 1:300 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234471 diluted at 1:300 and staining in paraffin-embedded human prostate tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: Colo320 whole cell lysate, Rat liver tissueAll lanes:

OLFM4 antibody at 10ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000

dilutionPredicted band size: 58 KDaObserved band size: 58 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-OLFM4 antibody	May promote proliferation of pancreatic cancer cells by favoring the transition from the S to G2/M phase. In myeloid leukemic cell lines, inhibits cell growth and induces cell differentiation and apoptosis. May play a role in the inhibition of EIF4EBP1 phosphorylation/deactivation. Facilitates cell adhesion, most probably through interaction with cell surface lectins and cadherin.
Product Categories/Family for anti-OLFM4 antibody	Cancer

NCBI AND UNIPROT INFORMATION

NCBI GI #	32313593
NCBI GeneID	10562
NCBI Accession #	NP_006409.3
NCBI GenBank Nucleotide #	NM_006418.4
UniProt Accession #	Q6UX06; O95362; Q5VWG0; Q86T22
Molecular Weight	57,280 Da
NCBI Official Full Name	olfactomedin-4
NCBI Official Synonym Full Names	olfactomedin 4
NCBI Official Symbol	OLFM4
NCBI Official Synonym Symbols	GC1; OLM4; OlfD; GW112; hGC-1; hOLfD; UNQ362; bA209J19.1
NCBI Protein Information	olfactomedin-4
UniProt Protein Name	Olfactomedin-4
UniProt Gene Name	OLFM4
UniProt Synonym Gene Names	GW112; OLM4; hGC-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/olfm4-antibody-immunohistochemistry-western-blot-elisa/234471>