

Rabbit anti-Human FOLR2 Polyclonal Antibody | anti-FOLR2 antibody

FOLR2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234773	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

FOLR2; Polyclonal Antibody; FOLR2 Antibody; Folate receptor beta; FR-beta; Folate receptor 2; Folate receptor; fetal/placental; Placental folate-binding protein; FBP; anti-FOLR2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	255
Applicable Applications for anti-FOLR2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Folate receptor beta protein (35-167AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells with AAA234773 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of AAA234773 diluted at 1:400 and staining in paraffin-embedded human placenta tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-FOLR2 antibody	Binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells. Has high affinity for folate and folic acid analogs at neutral pH. Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release.
Product Categories/Family for anti-FOLR2 antibody	Cancer; Metabolism; Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	166064050
NCBI GeneID	2350
NCBI Accession #	NP_000794.3
NCBI GenBank Nucleotide #	NM_000803.4
UniProt Accession #	P14207; Q05CA5; Q6GTE8
Molecular Weight	29,280 Da
NCBI Official Full Name	folate receptor beta
NCBI Official Synonym Full Names	folate receptor beta
NCBI Official Symbol	FOLR2
NCBI Official Synonym Symbols	FBP; FR-P3; FR-BETA; BETA-HFR; FBP/PL-1
NCBI Protein Information	folate receptor beta
UniProt Protein Name	Folate receptor beta
UniProt Gene Name	FOLR2
UniProt Synonym Gene Names	FR-beta; FBP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/folr2-antibody-immunofluorescence-immunohistochemistry-elisa/234773>