

Rabbit anti-Human OMD Polyclonal Antibody | anti-OMD antibody

OMD Antibody

CATALOG # AAA234970	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

OMD; Polyclonal Antibody; OMD Antibody; Osteomodulin; Keratan sulfate proteoglycan osteomodulin; KSPG osteomodulin; Osteoadherin; OSAD; SLRR2C; UNQ190/PRO216; anti-OMD antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	421
Applicable Applications for anti-OMD antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Osteomodulin protein (339-411AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image of AAA234970 diluted at 1:400 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234970 diluted at 1:400 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: A549 whole cell lysate, MCF-7 whole cell lysateAll lanes: OMD antibody at 3.5ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 50 KDaObserved band size: 50 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-OMD antibody	May be implicated in biomineralization processes. Has a function in binding of osteoblasts via the alpha(V)beta(3)-integrin (By similarity).
Product Categories/Family for anti-OMD antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	4826876
NCBI GeneID	4958
NCBI Accession #	NP_005005.1
NCBI GenBank Nucleotide #	NM_005014.2
UniProt Accession #	Q99983; Q5TBF4
Molecular Weight	49,492 Da
NCBI Official Full Name	osteomodulin
NCBI Official Synonym Full Names	osteomodulin
NCBI Official Symbol	OMD
NCBI Official Synonym Symbols	OSAD; SLRR2C
NCBI Protein Information	osteomodulin
UniProt Protein Name	Osteomodulin
UniProt Gene Name	OMD
UniProt Synonym Gene Names	SLRR2C; KSPG osteomodulin; OSAD

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/omd-antibody-immunohistochemistry-western-blot-elisa/234970>