

Rabbit anti-Human SSH1 Polyclonal Antibody | anti-SSH1 antibody

SSH1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235034	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

SSH1; Polyclonal Antibody; SSH1 Antibody; Protein phosphatase Slingshot homolog 1; SSH-like protein 1; SSH-1L; hSSH-1L; KIAA1298; SSH1L; anti-SSH1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	692
Applicable Applications for anti-SSH1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Protein phosphatase Slingshot homolog 1 protein (928-1049AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of Hela cells with AAA235034 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of AAA235034 diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-SSH1 antibody	Protein phosphatase which regulates actin filament dynamics. Dephosphorylates and activates the actin binding/depolymerizing factor cofilin, which subsequently binds to actin filaments and stimulates their disassembly. Inhibitory phosphorylation of cofilin is mediated by LIMK1, which may also be dephosphorylated and inactivated by this protein.
Product Categories/Family for anti-SSH1 antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	239047441
NCBI GeneID	54434
NCBI Accession #	NP_001154802.1
NCBI GenBank Nucleotide #	NM_001161330.1
UniProt Accession #	Q8WYL5; Q6P6C0; Q8N9A7; Q8WYL3; Q8WYL4; Q9P2P8
Molecular Weight	78,978 Da
NCBI Official Full Name	protein phosphatase Slingshot homolog 1 isoform 2
NCBI Official Synonym Full Names	slingshot protein phosphatase 1
NCBI Official Symbol	SSH1
NCBI Official Synonym Symbols	SSH1L
NCBI Protein Information	protein phosphatase Slingshot homolog 1
UniProt Protein Name	Protein phosphatase Slingshot homolog 1
UniProt Gene Name	SSH1
UniProt Synonym Gene Names	KIAA1298; SSH1L; SSH-1L; hSSH-1L

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ssh1-antibody-immunofluorescence-immunohistochemistry-elisa/235034>