

CD146 recombinant antibody

CD146 Antibody

CATALOG # AAA235164	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Western Blot, ELISA
-------------------------------	----------------------------	--

PRODUCT SYNONYMS

CD146; Monoclonal Recombinant Antibody; CD146 Antibody; Cell surface glycoprotein MUC18; Cell surface glycoprotein P1H12; Melanoma cell adhesion molecule; Melanoma-associated antigen A32; Melanoma-associated antigen MUC18; S-endo 1 endothelial-associated antigen; MCAM; MUC18; CD146 recombinant antibody

PRODUCT SPECIFICATIONS

Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	646
Applicable Applications for CD146 recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	A synthesized peptide
Conjugation	Non-conjugated
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



Application Data

Overlay histogram showing A375 cells stained with AAA235164 (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4 degree C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4 degree C. Control antibody (green line) was used under the same conditions. Acquisition of >10, 000 events was performed.



IHC (Immunohistochemisry)

Immunofluorescence staining of U251 cells with AAA235164 at 1:37, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of AAA235164 diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: A375 whole cell lysateAll lanes: CD146 antibody at 0.6ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 72 KDaObserved band size: 120 KDa

ADDITIONAL INFORMATION

Related Product Information for CD146 recombinant antibody	Plays a role in cell adhesion, and in cohesion of the endothelial monolayer at intercellular junctions in vascular tissue. Its expression may allow melanoma cells to interact with cellular elements of the vascular system, thereby enhancing hematogeneous tumor spread. Could be an adhesion molecule active in neural crest cells during embryonic development. Acts as surface receptor that triggers tyrosine phosphorylation of FYN and PTK2/FAK1, and a transient increase in the intracellular calcium concentration.
Product Categories/Family for CD146 recombinant antibody	Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	71274107
NCBI GeneID	4162
NCBI Accession #	NP_006491.2
NCBI GenBank Nucleotide #	NM_006500.2
UniProt Accession #	P43121; O95812; Q59E86; Q6PHR3; Q6ZTR2
Molecular Weight	57,605 Da
NCBI Official Full Name	cell surface glycoprotein MUC18
NCBI Official Synonym Full Names	melanoma cell adhesion molecule
NCBI Official Symbol	MCAM
NCBI Official Synonym Symbols	CD146; MUC18
NCBI Protein Information	cell surface glycoprotein MUC18
UniProt Protein Name	Cell surface glycoprotein MUC18
UniProt Gene Name	MCAM
UniProt Synonym Gene Names	MUC18

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd146-recombinant-antibody-flow-cytometry-immunofluorescence-immunohistochemistry-western-blot-elisa/235164>