

Histone H1.4 Recombinant Antibody | H1F4 recombinant antibody

Phospho-Histone H1.4 (T17) Antibody

CATALOG # AAA235171	REACTIVITY Human	APPLICATIONS Immunofluorescence, Immunohistochemistry, ELISA
-------------------------------	----------------------------	--


PRODUCT SYNONYMS

Histone H1.4; Monoclonal Recombinant Antibody; Phospho-Histone H1.4 (T17) Antibody; Histone H1b; Histone H1s-4; HIST1H1E; H1F4 recombinant antibody

PRODUCT SPECIFICATIONS


Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	219
Applicable Applications for H1F4 recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	A synthesized peptide
Conjugation	Non-conjugated
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescence staining of MCF-7 cells with AAA235171 at 1:56, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of AAA235171 diluted at 1:100 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA235171 diluted at 1:100 and staining in paraffin-embedded human lymph node tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for H1F4 recombinant antibody	Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation (By similarity).
Product Categories/Family for H1F4 recombinant antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885379
NCBI GeneID	3008
NCBI Accession #	NP_005312.1
NCBI GenBank Nucleotide #	NM_005321.2
UniProt Accession #	P10412; Q4VB25
Molecular Weight	21,865 Da
NCBI Official Full Name	histone H1.4
NCBI Official Synonym Full Names	histone cluster 1 H1 family member e
NCBI Official Symbol	HIST1H1E
NCBI Official Synonym Symbols	H1E; H1.4; H1F4; RMNS; H1s-4; dJ221C16.5
NCBI Protein Information	histone H1.4
UniProt Protein Name	Histone H1.4
UniProt Gene Name	HIST1H1E
UniProt Synonym Gene Names	H1F4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/histone-h14-recombinant-antibody-immunofluorescence-immunohistochemistry-elisa/235171>