

Mono-methyl-Histone H3.1 Recombinant Antibody | HIST1H3A recombinant antibody

Mono-methyl-Histone H3.1 (R2) Antibody

CATALOG # AAA235174	REACTIVITY Human	APPLICATIONS Immunofluorescence, Immunocytochemistry, Western Blot, ELISA
-------------------------------	----------------------------	---


PRODUCT SYNONYMS

Mono-methyl-Histone H3.1; Monoclonal Recombinant Antibody; Mono-methyl-Histone H3.1 (R2) Antibody; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/h; Histone H3/i; Histone H3/j; Histone H3/k; Histone H3/l; H3FA; AND; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; HIST1H3A recombinant antibody

PRODUCT SPECIFICATIONS


Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	136
Applicable Applications for HIST1H3A recombinant antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot), ELISA
Immunogen	A synthesized peptide
Conjugation	Non-conjugated
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescence staining of HeLa cells with AAA235174 at 1:60, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



WB (Western Blot)

Western Blot Positive WB detected in: HeLa whole cell lysate, HepG2 whole cell lysate, SH-SY5Y whole cell lysate All lanes: Mono-methyl-Histone H3.1 (R2) antibody at 0.95ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 15 KDa Observed band size: 15 KDa



ICC (Immunocytochemistry)

Immunocytochemistry analysis of AAA235174 diluted at 1:100 and staining in HeLa cells performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for HIST1H3A recombinant antibody	Epigenetics and Nuclear Signaling
---	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504281
NCBI GeneID	8350
NCBI Accession #	NP_003520.1
NCBI GenBank Nucleotide #	NM_003529.2
UniProt Accession #	P68431; P02295; P02296; P16106; Q6ISV8; Q6NWP8; Q6NWP9; Q6NXU4; Q71DJ3; Q93081; A0PJT7; A5PLR1
Molecular Weight	15,404 Da
NCBI Official Full Name	histone H3.1
NCBI Official Synonym Full Names	histone cluster 1 H3 family member a
NCBI Official Symbol	HIST1H3A
NCBI Official Synonym Symbols	H3/A; H3FA
NCBI Protein Information	histone H3.1
UniProt Protein Name	Histone H3.1
UniProt Gene Name	HIST1H3A
UniProt Synonym Gene Names	H3FA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/mono-methyl-histone-h31-recombinant-antibody-immunofluorescence-immunocytochemistry-western-blot-elisa/235174>