

CD31 recombinant antibody

CD31 Antibody

CATALOG # AAA235180	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot, ELISA
-------------------------------	----------------------------	--

PRODUCT SYNONYMS

CD31; Monoclonal Recombinant Antibody; CD31 Antibody; Platelet endothelial cell adhesion molecule; PECAM-1; EndoCAM; GPIIA⁺; PECA1; PECAM1; CD31 recombinant antibody

PRODUCT SPECIFICATIONS


Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	738
Applicable Applications for CD31 recombinant antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	A synthesized peptide
Conjugation	Non-conjugated
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




Application Data

Overlay histogram showing Jurkat cells stained with AAA235180 (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4 degree C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4 degree C. Control antibody (green line) was used under the same conditions. Acquisition of >10, 000 events was performed.



IHC (Immunohistochemistry)

IHC image of AAA235180 diluted at 1:100 and staining in paraffin-embedded human tonsil tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: THP-1 whole cell lysate All lanes: CD31 antibody at 0.95ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 125 KDa Observed band size: 125 KDa

ADDITIONAL INFORMATION

Related Product Information for CD31 recombinant antibody	Induces susceptibility to atherosclerosis (By similarity). Cell adhesion molecule which is required for leukocyte transendothelial migration (TEM) under most inflammatory conditions. Tyr-690 plays a critical role in TEM and is required for efficient trafficking of PECAM1 to and from the lateral border recycling compartment (LBRC) and is also essential for the LBRC membrane to be targeted around migrating leukocytes. Prevents phagocyte ingestion of closely apposed viable cells by transmitting 'detachment' signals, and changes function on apoptosis, promoting tethering of dying cells to phagocytes (the encounter of a viable cell with a phagocyte via the homophilic interaction of PECAM1 on both cell surfaces leads to the viable cell's active repulsion from the phagocyte. During apoptosis, the inside-out signaling of PECAM1 is somehow disabled so that the apoptotic cell does not actively reject the phagocyte anymore. The lack of this repulsion signal together with the interaction of the eat-me signals and their respective receptors causes the attachment of the apoptotic cell to the phagocyte, thus triggering the process of engulfment). Isoform Delta15 is unable to protect against apoptosis. Modulates BDKRB2 activation. Regulates bradykinin- and hyperosmotic shock-induced ERK1/2 activation in human umbilical cord vein cells (HUVEC).
Product Categories/Family for CD31 recombinant antibody	Cardiovascular

NCBI AND UNIPROT INFORMATION

NCBI GI #	313760624
NCBI GeneID	5175
NCBI Accession #	NP_000433.4
NCBI GenBank Nucleotide #	NM_000442.4
UniProt Accession #	P16284; Q6LDA9; Q8TBH1; Q96RF5; Q96RF6; Q9NP65; Q9NPB7; Q9NPG9; Q9NQS9; Q9NQT0; A8K3S7; D3DU31
Molecular Weight	81,378 Da
NCBI Official Full Name	platelet endothelial cell adhesion molecule
NCBI Official Synonym Full Names	platelet and endothelial cell adhesion molecule 1
NCBI Official Symbol	PECAM1
NCBI Official Synonym Symbols	CD31; PECA1; GPIIA'; PECAM-1; endoCAM; CD31/EndoCAM
NCBI Protein Information	platelet endothelial cell adhesion molecule
UniProt Protein Name	Platelet endothelial cell adhesion molecule
UniProt Gene Name	PECAM1
UniProt Synonym Gene Names	PECAM-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd31-recombinant-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/235180>