

Histone H3.1 Recombinant Antibody | HIST1H3A recombinant antibody

Histone H3.1 Antibody

CATALOG # AAA235190	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Western Blot, ELISA
-------------------------------	----------------------------	--

PRODUCT SYNONYMS

Histone H3.1; Monoclonal Recombinant Antibody; Histone H3.1 Antibody; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/h; Histone H3/i; Histone H3/j; Histone H3/k; Histone H3/l; H3FA; AND; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; HIST1H3A recombinant antibody

PRODUCT SPECIFICATIONS


Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	136
Applicable Applications for HIST1H3A recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	A synthesized peptide
Conjugation	Non-conjugated
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



Application Data

Overlay histogram showing HeLa cells stained with AAA235190 (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4 degree C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4 degree C. Control antibody (green line) was used under the same conditions. Acquisition of >10, 000 events was performed.



IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells with AAA235190 at 1:93, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).




IHC (Immunohistochemisry)

IHC image of AAA235190 diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohiostchemistry)

IHC image of AAA235190 diluted at 1:100 and staining in paraffin-embedded human glioma cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: Jurkat whole cell lysateAll lanes: Histone H3.1 antibody at 1.5ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 15 KDaObserved band size: 15 KDa

ADDITIONAL INFORMATION

Related Product Information for HIST1H3A recombinant antibody	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Product Categories/Family for HIST1H3A recombinant antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504281
NCBI GeneID	8350
NCBI Accession #	NP_003520.1
NCBI GenBank Nucleotide #	NM_003529.2
UniProt Accession #	P68431 ; P02295 ; P02296 ; P16106 ; Q6ISV8 ; Q6NWP8 ; Q6NWP9 ; Q6NXU4 ; Q71DJ3 ; Q93081 ; A0PJT7 ; A5PLR1
Molecular Weight	15,404 Da
NCBI Official Full Name	histone H3.1
NCBI Official Synonym Full Names	histone cluster 1 H3 family member a
NCBI Official Symbol	HIST1H3A
NCBI Official Synonym Symbols	H3/A; H3FA
NCBI Protein Information	histone H3.1
UniProt Protein Name	Histone H3.1
UniProt Gene Name	HIST1H3A
UniProt Synonym Gene Names	H3FA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.