

CD99 recombinant antibody

CD99 Antibody

CATALOG # AAA235191	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot, ELISA
-------------------------------	----------------------------	--

PRODUCT SYNONYMS

CD99; Monoclonal Recombinant Antibody; CD99 Antibody; CD99 antigen; 12E7; E2 antigen; Protein MIC2; T-cell surface glycoprotein E2; MIC2; MIC2X; MIC2Y; CD99 recombinant antibody

PRODUCT SPECIFICATIONS

Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	169
Applicable Applications for CD99 recombinant antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	A synthesized peptide
Conjugation	Non-conjugated
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




Application Data

Overlay histogram showing Jurkat cells stained with AAA235191 (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4 degree C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4 degree C. Control antibody (green line) was used under the same conditions. Acquisition of >10, 000 events was performed.



IHC (Immunohistochemistry)

IHC image of AAA235191 diluted at 1:100 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: Jurkat whole cell lysate All lanes: CD99 antibody at 0.8ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 28 KDa Observed band size: 28 KDa

ADDITIONAL INFORMATION

Related Product Information for CD99 recombinant antibody	Involved in T-cell adhesion processes and in spontaneous rosette formation with erythrocytes. Plays a role in a late step of leukocyte extravasation helping leukocytes to overcome the endothelial basement membrane. Acts at the same site as, but independently of, PECAM1. Involved in T-cell adhesion processes (By similarity).
Product Categories/Family for CD99 recombinant antibody	Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	171543879
NCBI GeneID	4267
NCBI Accession #	NP_001116370.1
NCBI GenBank Nucleotide #	NM_001122898.2
UniProt Accession #	P14209; O00518; Q6ICV7; A6NIW1
Molecular Weight	17,128 Da
NCBI Official Full Name	CD99 antigen isoform b
NCBI Official Synonym Full Names	CD99 molecule (Xg blood group)
NCBI Official Symbol	CD99
NCBI Official Synonym Symbols	MIC2; HBA71; MIC2X; MIC2Y; MSK5X
NCBI Protein Information	CD99 antigen
UniProt Protein Name	CD99 antigen
UniProt Gene Name	CD99
UniProt Synonym Gene Names	MIC2; MIC2X; MIC2Y

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd99-recombinant-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/235191>