

Rabbit anti-Human CYP21A2 Polyclonal Antibody | anti-CYP21A2 antibody

CYP21A2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235292	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

CYP21A2; Polyclonal Antibody; CYP21A2 Antibody; Steroid 21-hydroxylase; EC 1.14.14.16; 21-OHase; Cytochrome P-450c21; Cytochrome P450 21; Cytochrome P450 XXI; Cytochrome P450-C21; Cytochrome P450-C21B; CYP21 CYP21B; anti-CYP21A2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95% Protein G Purified
Form/Format	Liquid
Sequence Length	495
Applicable Applications for anti-CYP21A2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Steroid 21-hydroxylase protein (312-418AA)
Conjugate	Non-Conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HepG2 with CSB-PA006400LA01HU at 1:33, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of CSB-PA006400LA01HU diluted at 1:100 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: Jurkat whole cell lysate, HL-60 whole cell lysate, 293 whole cell lysate, SH-SY5Y whole cell lysate, Hela whole cell lysate, PC3 whole cell lysateAll lanes: CYP21A2 antibody at 1:500SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 56, 53 kDaObserved band size: 56 kDa

ADDITIONAL INFORMATION

Product Categories/Family for anti-CYP21A2 antibody	Cardiovascular
---	----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	323510663
NCBI GeneID	1589
NCBI Accession #	NP_000491.4
NCBI GenBank Nucleotide #	NM_000500.9
UniProt Accession #	P08686; P04033; Q01204; Q08AG8; Q16749; Q16806; Q5ST44; Q96NU8; A2BHY6
Molecular Weight	52,597 Da
NCBI Official Full Name	steroid 21-hydroxylase isoform a
NCBI Official Synonym Full Names	cytochrome P450 family 21 subfamily A member 2
NCBI Official Symbol	CYP21A2
NCBI Official Synonym Symbols	CAH1; CPS1; CA21H; CYP21; CYP21B; P450c21B
NCBI Protein Information	steroid 21-hydroxylase
UniProt Protein Name	Steroid 21-hydroxylase
UniProt Gene Name	CYP21A2
UniProt Synonym Gene Names	CYP21; CYP21B

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyp21a2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/235292>