

Rabbit anti-Human, Rat CABIN1 Polyclonal Antibody | anti-CABIN1 antibody

CABIN1 Antibody

CATALOG # AAA235388	HOST Rabbit	REACTIVITY Human, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	---------------------------------	--

PRODUCT SYNONYMS

CABIN1; Polyclonal Antibody; CABIN1 Antibody; Calcineurin-binding protein cabin-1; Calcineurin inhibitor; CAIN; KIAA0330; anti-CABIN1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95% Protein G Purified
Form/Format	Liquid
Sequence Length	2220
Applicable Applications for anti-CABIN1 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Calcineurin-binding protein cabin-1 protein (1978-2123AA)
Conjugate	Non-Conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



WB (Western Blot)

Western Blot Positive WB detected in: A549 whole cell lysate, Rat brain tissue All lanes: CABIN1 antibody at 3.1ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 247, 238 kDa Observed band size: 247 kDa



IHC (Immunohistochemistry)

IHC image of CSB-PA23399A0Rb diluted at 1:400 and staining in paraffin-embedded human cervical cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of CSB-PA23399A0Rb diluted at 1:400 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CABIN1 antibody	Signal Transduction
--	---------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	313151181
NCBI GeneID	23523
NCBI Accession #	NP_001186210.1
NCBI GenBank Nucleotide #	NM_001199281.1
UniProt Accession #	Q9Y6J0; Q6PHY0; Q9Y460; G5E9F3
Molecular Weight	237,574 Da
NCBI Official Full Name	calcineurin-binding protein cabin-1 isoform a
NCBI Official Synonym Full Names	calcineurin binding protein 1
NCBI Official Symbol	CABIN1
NCBI Official Synonym Symbols	CAIN; PPP3IN; KB-318B8.7
NCBI Protein Information	calcineurin-binding protein cabin-1
UniProt Protein Name	Calcineurin-binding protein cabin-1
UniProt Gene Name	CABIN1
UniProt Synonym Gene Names	KIAA0330; CAIN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cabin1-antibody-immunohistochemistry-western-blot-elisa/235388>