

ATF2 recombinant antibody

ATF2 Antibody

CATALOG # AAA235522	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

ATF2; Monoclonal Recombinant Antibody; ATF2 Antibody; Cyclic AMP-dependent transcription factor ATF-2; Activating transcription factor 2; Cyclic AMP-responsive element-binding protein 2; CREB-2; cAMP-responsive element-binding protein 2; HB16; Histone acetyltransferase ATF2; cAMP response element-binding protein CRE-BP1; CREB2; CREBP1; ATF2 recombinant antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	505
Applicable Applications for ATF2 recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human ATF2
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of CSB-RA002270A0HU diluted at 1:115.5 and staining in paraffin-embedded human prostate tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: K562 whole cell lysate, 293T whole cell lysate All lanes: ATF2 antibody at 1.2ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 55, 36, 24, 49, 53, 14, 25, 16 kDa Observed band size: 70 kDa

ADDITIONAL INFORMATION

Related Product Information for ATF2 recombinant antibody	Transcriptional activator which regulates the transcription of various genes, including those involved in anti-apoptosis, cell growth, and DNA damage response. Dependent on its binding partner, binds to CRE (cAMP response element) consensus sequences (5'-TGACGTCA-3') or to AP-1 (activator protein 1) consensus sequences (5'-TGACTCA-3'). In the nucleus, contributes to global transcription and the DNA damage response, in addition to specific transcriptional activities that are related to cell development, proliferation and death. In the cytoplasm, interacts with and perturbs HK1-and VDAC1-containing complexes at the mitochondrial outer membrane, thereby impairing mitochondrial membrane potential, inducing mitochondrial leakage and promoting cell death. The phosphorylated form (mediated by ATM) plays a role in the DNA damage response and is involved in the ionizing radiation (IR)-induced S phase checkpoint control and in the recruitment of the MRN complex into the IR-induced foci (IRIF). Exhibits histone acetyltransferase (HAT) activity which specifically acetylates histones H2B and H4 in vitro. In concert with CUL3 and RBX1, promotes the degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM. Can elicit oncogenic or tumor suppressor activities depending on the tissue or cell type.
Product Categories/Family for ATF2 recombinant antibody	Recombinant Antibody; Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	368711269
NCBI GeneID	1386
NCBI Accession #	NP_001243019.1
NCBI GenBank Nucleotide #	NM_001256090.1
UniProt Accession #	P15336; Q13000; Q3B7B7; Q4ZFU9; Q53RY2; Q8TAR1; A1L3Z2; A4D7U4; A4D7U5; A4D7V1; D3DPE9; G8JLM5
Molecular Weight	35,044 Da
NCBI Official Full Name	cyclic AMP-dependent transcription factor ATF-2 isoform 1
NCBI Official Synonym Full Names	activating transcription factor 2
NCBI Official Symbol	ATF2
NCBI Official Synonym Symbols	HB16; CREB2; TREB7; CREB-2; CRE-BP1
NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-2
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-2
UniProt Gene Name	ATF2
UniProt Synonym Gene Names	CREB2; CREBP1; cAMP-dependent transcription factor ATF-2; CREB-2; cAMP-responsive element-binding protein 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/atf2-recombinant-antibody-immunohistochemistry-western-blot-elisa/235522>