

BAG3 recombinant antibody

BAG3 Antibody

CATALOG # AAA235525	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

BAG3; Monoclonal Recombinant Antibody; BAG3 Antibody; BAG family molecular chaperone regulator 3; BAG-3; Bcl-2-associated athanogene 3; Bcl-2-binding protein Bis; Docking protein CAIR-1; BIS; BAG3 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	575
Applicable Applications for BAG3 recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human BAG3
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of CSB-RA002531A0HU diluted at 1:73.75 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: HeLa whole cell lysate, 293 whole cell lysate, A549 whole cell lysate, HepG2 whole cell lysate, U87 whole cell lysate. All lanes: BAG3 antibody at 0.73ug/ml. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 62 KDa. Observed band size: 80 KDa.

ADDITIONAL INFORMATION

Related Product Information for BAG3 recombinant antibody	Co-chaperone for HSP70 and HSC70 chaperone proteins. Acts as a nucleotide-exchange factor (NEF) promoting the release of ADP from the HSP70 and HSC70 proteins thereby triggering client/substrate protein release. Nucleotide release is mediated via its binding to the nucleotide-binding domain (NBD) of HSPA8/HSC70 where as the substrate release is mediated via its binding to the substrate-binding domain (SBD) of HSPA8/HSC70 (PubMed:9873016, PubMed:27474739). Has anti-apoptotic activity (PubMed:10597216). Plays a role in the HSF1 nucleocytoplasmic transport (PubMed:26159920).
Product Categories/Family for BAG3 recombinant antibody	Recombinant Antibody; Cell Biology

NCBI AND UNIPROT INFORMATION

NCBI GI #	14043024
NCBI GeneID	9531
NCBI Accession #	NP_004272.2
NCBI GenBank Nucleotide #	NM_004281.3
UniProt Accession #	O95817; Q3B763; Q9NT20; Q9P120; A8K5L8
Molecular Weight	61,595 Da
NCBI Official Full Name	BAG family molecular chaperone regulator 3
NCBI Official Synonym Full Names	BCL2 associated athanogene 3
NCBI Official Symbol	BAG3
NCBI Official Synonym Symbols	BIS; MFM6; BAG-3; CAIR-1
NCBI Protein Information	BAG family molecular chaperone regulator 3
UniProt Protein Name	BAG family molecular chaperone regulator 3
UniProt Gene Name	BAG3
UniProt Synonym Gene Names	BIS; BAG-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/bag3-recombinant-antibody-immunohistochemistry-western-blot-elisa/235525>