

CREB1 recombinant antibody

Phospho-CREB1 (S133) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235534	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

CREB1; Monoclonal Recombinant Antibody; Phospho-CREB1 (S133) Antibody; Cyclic AMP-responsive element-binding protein 1; CREB-1; cAMP-responsive element-binding protein 1; CREB1 recombinant antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	327
Applicable Applications for CREB1 recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human Phospho-CREB1 (S133)
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells with CSB-RA005947A133phHU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of CSB-RA005947A133phHU diluted at 1:100 and staining in paraffin-embedded human lung cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of CSB-RA005947A133pHU diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in?Hela whole cell lysate,293 whole cell lysateAll lanes?Phospho-CREB1 antibody at 1.65ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 46 KDaObserved band size: 46 KDa

ADDITIONAL INFORMATION

Related Product Information for CREB1 recombinant antibody	Phosphorylation-dependent transcription factor that stimulates transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells.
Product Categories/Family for CREB1 recombinant antibody	Recombinant Antibody ; Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758054
NCBI GeneID	1385
NCBI Accession #	NP_004370.1
NCBI GenBank Nucleotide #	NM_004379.5
UniProt Accession #	P16220 ; P21934 ; Q6V963 ; Q9UMA7
Molecular Weight	35,136 Da
NCBI Official Full Name	cyclic AMP-responsive element-binding protein 1 isoform A
NCBI Official Synonym Full Names	cAMP responsive element binding protein 1
NCBI Official Symbol	CREB1
NCBI Official Synonym Symbols	CREB; CREB-1
NCBI Protein Information	cyclic AMP-responsive element-binding protein 1
UniProt Protein Name	Cyclic AMP-responsive element-binding protein 1
UniProt Gene Name	CREB1
UniProt Synonym Gene Names	CREB-1; cAMP-responsive element-binding protein 1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/creb1-recombinant-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/235534>