

FEN1 recombinant antibody

FEN1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235540	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

FEN1; Monoclonal Recombinant Antibody; FEN1 Antibody; Flap endonuclease 1UniRule annotation; DNase IV; Flap structure-specific endonuclease 1UniRule annotation; FEN1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	380
Applicable Applications for FEN1 recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human FEN1
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells with CSB-RA008585A0HU at 1:25, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of CSB-RA008585A0HU diluted at 1:77.5 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: HeLa whole cell lysate, Raji whole cell lysate, HepG2 whole cell lysate, Jurkat whole cell lysate, MCF-7 whole cell lysateAll lanes: FEN1 antibody at 0.775ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 43, 36 KDaObserved band size: 43 KDa

ADDITIONAL INFORMATION

Related Product Information for FEN1 recombinant antibody	Structure-specific nuclease with 5'-flap endonuclease and 5'-3' exonuclease activities involved in DNA replication and repair. During DNA replication, cleaves the 5'-overhanging flap structure that is generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. It enters the flap from the 5'-end and then tracks to cleave the flap base, leaving a nick for ligation. Also involved in the long patch base excision repair (LP-BER) pathway, by cleaving within the apurinic/aprimidinic (AP) site-terminated flap. Acts as a genome stabilization factor that prevents flaps from equilibrating into structures that lead to duplications and deletions. Also possesses 5'-3' exonuclease activity on nicked or gapped double-stranded DNA, and exhibits RNase H activity. Also involved in replication and repair of rDNA and in repairing mitochondrial DNA.
Product Categories/Family for FEN1 recombinant antibody	Recombinant Antibody ; Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758356
NCBI GeneID	2237
NCBI Accession #	NP_004102.1
NCBI GenBank Nucleotide #	NM_004111.6
UniProt Accession #	P39748
Molecular Weight	35,673 Da
NCBI Official Full Name	flap endonuclease 1
NCBI Official Synonym Full Names	flap structure-specific endonuclease 1
NCBI Official Symbol	FEN1
NCBI Official Synonym Symbols	MF1; RAD2; FEN-1
NCBI Protein Information	flap endonuclease 1
UniProt Protein Name	Flap endonuclease 1
UniProt Gene Name	FEN1
UniProt Synonym Gene Names	RAD2; FEN-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/fen1-recombinant-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/235540>