

JUN recombinant antibody

Phospho-JUN (S63) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235568	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

JUN; Monoclonal Recombinant Antibody; Phospho-JUN (S63) Antibody; Transcription factor AP-1; Activator protein 1; AP1; Proto-oncogene c-Jun; V-jun avian sarcoma virus 17 oncogene homolog; p39; JUN recombinant antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	331
Applicable Applications for JUN recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human Phospho-JUN (S63)
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of A549 cells with CSB-RA011972A63pHU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of CSB-RA011972A63pHU diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in?Hela whole cell lysate,A549 whole cell lysate(treated with Calyculin A or EGF)All lanes?Phospho-JUN antibody at 0.95ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 48 KDaObserved band size: 48 KDa

ADDITIONAL INFORMATION

Related Product Information for JUN recombinant antibody	Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation. Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306).
Product Categories/Family for JUN recombinant antibody	Recombinant Antibody; Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758616
NCBI GeneID	3725
NCBI Accession #	NP_002219.1
NCBI GenBank Nucleotide #	NM_002228.3
UniProt Accession #	P05412; Q6FHM7; Q96G93
Molecular Weight	35,676 Da
NCBI Official Full Name	transcription factor AP-1
NCBI Official Synonym Full Names	Jun proto-oncogene, AP-1 transcription factor subunit
NCBI Official Symbol	JUN
NCBI Official Synonym Symbols	AP1; p39; AP-1; c-Jun
NCBI Protein Information	transcription factor AP-1
UniProt Protein Name	Transcription factor AP-1
UniProt Gene Name	JUN
UniProt Synonym Gene Names	AP1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/jun-recombinant-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/235568>