

RPS6KB1 recombinant antibody

Phospho-RPS6KB1 (T421+S424) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235587	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Western Blot, ELISA


PRODUCT SYNONYMS

RPS6KB1; Monoclonal Recombinant Antibody; Phospho-RPS6KB1 (T421+S424) Antibody; Ribosomal protein S6 kinase beta-1; 70 kDa ribosomal protein S6 kinase 1; P70S6K1; p70-S6K 1; Ribosomal protein S6 kinase I; Serine/threonine-protein kinase 14A; p70 ribosomal S6 kinase alpha; p70 S6 kinase alpha; p70 S6K-alpha; p70 S6KA; STK14A; RPS6KB1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	502
Applicable Applications for RPS6KB1 recombinant antibody	IP (Immunoprecipitation), IF (Immunofluorescence), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human Phospho-RPS6KB1 (T421+S424)
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IP (Immunoprecipitation)

Immunoprecipitating Phospho-RPS6KB1 in HeLa whole cell lysate Lane 1: Rabbit control IgG(1ug) instead of CSB-RA020470A421phHU in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000) Lane 2: CSB-RA020470A421phHU(3ug)+ HeLa whole cell lysate(1mg) Lane 3: HeLa whole cell lysate (20ug)



WB (Western Blot)

Western Blot Positive WB detected in 293 whole cell lysate (treated with Calyculin A or not) All lanes? Phospho-RPS6KB1 antibody at 0.93ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 70 kDa Observed band size: 70 kDa



IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells with CSB-RA020470A421phHU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).

ADDITIONAL INFORMATION

Related Product Information for RPS6KB1 recombinant antibody	Serine/threonine-protein kinase that acts downstream of mTOR signaling in response to growth factors and nutrients to promote cell proliferation, cell growth and cell cycle progression. Regulates protein synthesis through phosphorylation of EIF4B, RPS6 and EEF2K, and contributes to cell survival by repressing the pro-apoptotic function of BAD. Under conditions of nutrient depletion, the inactive form associates with the EIF3 translation initiation complex. Upon mitogenic stimulation, phosphorylation by the mammalian target of rapamycin complex 1 (mTORC1) leads to dissociation from the EIF3 complex and activation. The active form then phosphorylates and activates several substrates in the pre-initiation complex, including the EIF2B complex and the cap-binding complex component EIF4B. Also controls translation initiation by phosphorylating a negative regulator of EIF4A, PDCD4, targeting it for ubiquitination and subsequent proteolysis. Promotes initiation of the pioneer round of protein synthesis by phosphorylating POLDIP3/SKAR. In response to IGF1, activates translation elongation by phosphorylating EEF2 kinase (EEF2K), which leads to its inhibition and thus activation of EEF2. Also plays a role in feedback regulation of mTORC2 by mTORC1 by phosphorylating RICTOR, resulting in the inhibition of mTORC2 and AKT1 signaling. Mediates cell survival by phosphorylating the pro-apoptotic protein BAD and suppressing its pro-apoptotic function. Phosphorylates mitochondrial URI1 leading to dissociation of a URI1-PPP1CC complex. The free mitochondrial PPP1CC can then dephosphorylate RPS6KB1 at Thr-412, which is proposed to be a negative feedback mechanism for the RPS6KB1 anti-apoptotic function. Mediates TNF-alpha-induced insulin resistance by phosphorylating IRS1 at multiple serine residues, resulting in accelerated degradation of IRS1. In cells lacking functional TSC1-2 complex, constitutively phosphorylates and inhibits GSK3B. May be involved in cytoskeletal rearrangement through binding to neurabin. Phosphorylates and activates the pyrimidine biosynthesis enzyme CAD, downstream of MTOR.
Product Categories/Family for RPS6KB1 recombinant antibody	Recombinant Antibody; Cell Biology

NCBI AND UNIPROT INFORMATION

NCBI GI #	440546395
NCBI GeneID	6198
NCBI Accession #	NP_001258971.1
NCBI GenBank Nucleotide #	NM_001272042.1
UniProt Accession #	P23443; Q7Z721; B2R779; B4DLT4; B4DTG1; E7ESB8; F6UYM1
Molecular Weight	56,189 Da
NCBI Official Full Name	ribosomal protein S6 kinase beta-1 isoform b
NCBI Official Synonym Full Names	ribosomal protein S6 kinase B1
NCBI Official Symbol	RPS6KB1
NCBI Official Synonym Symbols	S6K; PS6K; S6K1; STK14A; p70-S6K; p70 S6KA; p70-alpha; S6K-beta-1; p70(S6K)-alpha
NCBI Protein Information	ribosomal protein S6 kinase beta-1
UniProt Protein Name	Ribosomal protein S6 kinase beta-1
UniProt Gene Name	RPS6KB1
UniProt Synonym Gene Names	STK14A; S6K-beta-1; S6K1; P70S6K1; p70-S6K 1; p70 S6 kinase alpha; p70 S6K-alpha; p70 S6KA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/rps6kb1-recombinant-antibody-immunoprecipitation-immunofluorescence-western-blot-elisa/235587>