

Mouse anti-Human Sarcomeric Actinin Alpha 2/ACTN2 Monoclonal Antibody | anti-ACTN2 antibody

Sarcomeric Actinin Alpha 2/ACTN2


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA23944	Mouse	Human	Western Blot, Immunohistochemistry

PRODUCT SYNONYMS

Sarcomeric Actinin Alpha 2/ACTN2; Monoclonal Antibody; Actin binding protein; Alpha actinin skeletal muscle; isoform 2; CMD1AA; F-actin cross-linking protein; anti-ACTN2 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG2b, kappa
Clone Number	ACTN2/3295
Specificity	Alpha actinins belong to the spectrin gene superfamily which represents a diverse group of cytoskeletal proteins, including the alpha and beta spectrins and dystrophins. Alpha actinin is an actin-binding protein with multiple roles in different cell types. In non-muscle cells, the cytoskeletal isoform is found along microfilament bundles and adherens-type junctions, where it is involved in binding actin to the membrane. In contrast, skeletal, cardiac, and smooth muscle isoforms are localized to the Z-disc and analogous dense bodies, where they help anchor the myofibrillar actin filaments. This gene encodes a muscle-specific, alpha actinin isoform that is expressed in both skeletal and cardiac muscles. Several transcript variants encoding different isoforms have been found for this gene.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	686
Applicable Applications for anti-ACTN2 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Application Notes	WB: 1-2ug/ml Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)
Positive Control	Skeletal or Cardiac Muscle.
Immunogen	A recombinant fragment (aa557-692) of human ACTN2 protein (exact sequence is proprietary)
Cellular Localization	Cytoplasmic
Hu-chromosome Location	1q43
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C. Antibody without azide: store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.




Application Data


Analysis of Protein Array containing more than 19,000 full-length human proteins using Sarcomeric Actinin Alpha 2 Mouse Monoclonal Antibody (ACTN2/3295). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.




WB (Western Blot)
Western Blot of human Skeletal muscle tissue lysates using Sarcomeric Actinin Alpha 2 Mouse Monoclonal Antibody (ACTN2/3295).




SDS-PAGE
SDS-PAGE Analysis Purified ACTN2 Mouse Monoclonal Antibody (ACTN2/3295). Confirmation of Purity and Integrity of Antibody.



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Cardiac Muscle stained with Sarcomeric Actinin Alpha 2 Mouse Monoclonal Antibody (ACTN2/3295).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Skeletal Muscle stained with Sarcomeric Actinin Alpha 2 Mouse Monoclonal Antibody (ACTN2/3295).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Skeletal Muscle stained with Sarcomeric Actinin Alpha 2 Mouse Monoclonal Antibody (ACTN2/3295).

ADDITIONAL INFORMATION

References	Baris, O., et al. Oncogene 24 (April 4, 2005): 4155.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	507588250
-----------	-----------

NCBI GeneID	88
NCBI Accession #	NP_001265273.1
NCBI GenBank Nucleotide #	NP_001265273.1
UniProt Accession #	P35609
Molecular Weight	103kDa
NCBI Official Full Name	alpha-actinin-2 isoform 3
NCBI Official Synonym Full Names	actinin alpha 2
NCBI Official Symbol	ACTN2
NCBI Official Synonym Symbols	CMH23; CMD1AA
NCBI Protein Information	alpha-actinin-2
UniProt Protein Name	Alpha-actinin-2
UniProt Gene Name	ACTN2
UniProt Entry Name	ACTN2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/sarcomeric-actinin-alpha-2-actn2-antibody-clone-actn2-3295-western-blot-immunohistochemistry-elisa/23944>