

Rabbit anti-Human EIF2AK3 Polyclonal Antibody | anti-EIF2AK3 antibody

EIF2AK3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243424	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

EIF2AK3; Polyclonal Antibody; EIF2AK3 Antibody; Eukaryotic Translation initiation factor 2-alpha kinase 3 (EC 2.7.11.1) (PRKR-like endoplasmic reticulum kinase) (Pancreatic eIF2-alpha kinase) (HsPEK); PEK PERK; anti-EIF2AK3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G Purified
Form/Format	Liquid Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Applicable Applications for anti-EIF2AK3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Eukaryotic translation initiation factor 2-alpha kinase 3 protein (115-330AA)
Immunogen Species	Human
Conjugate	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells with AAA243424 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of AAA243424 diluted at 1:300 and staining in paraffin-embedded human small intestine tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA243424 diluted at 1:300 and staining in paraffin-embedded human ovarian cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-EIF2AK3 antibody

Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	134304838
NCBI GeneID	9451
NCBI Accession #	NP_004827.4
NCBI GenBank Nucleotide #	NM_004836.5
UniProt Accession #	Q9NZJ5; O95846; Q53QY0; Q53SB1; A0AVH1; A0AVH2; B2RCU9
Molecular Weight	125,216 Da
NCBI Official Full Name	eukaryotic translation initiation factor 2-alpha kinase 3
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 2-alpha kinase 3
NCBI Official Symbol	EIF2AK3
NCBI Official Synonym Symbols	PEK; WRS; PERK
NCBI Protein Information	eukaryotic translation initiation factor 2-alpha kinase 3; hsPEK; pancreatic EIF2-alpha kinase; PRKR-like endoplasmic reticulum kinase; eukaryotic translation initiation factor 2 alpha kinase 3
UniProt Protein Name	Eukaryotic translation initiation factor 2-alpha kinase 3
UniProt Gene Name	EIF2AK3
UniProt Synonym Gene Names	PEK; PERK; HsPEK
UniProt Entry Name	E2AK3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/EIF2ak3-antibody-immunofluorescence-immunohistochemistry-elisa/243424>