

Rabbit anti-Human INSRR Polyclonal Antibody | anti-INSRR antibody

INSRR Antibody

CATALOG # AAA243554	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

INSRR; Polyclonal Antibody; INSRR Antibody; Insulin receptor-related protein (IRR) (EC 2.7.10.1) (IR-related receptor) [Cleaved into: Insulin receptor-related protein alpha chain; Insulin receptor-related protein beta chain]; IRR; anti-INSRR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Antigen Affinity Purified & Affinity Purified
Form/Format	Liquid Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Applicable Applications for anti-INSRR antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Insulin receptor-related protein (747-921AA)
Immunogen Species	Human
Conjugate	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image of AAA243554 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA243554 diluted at 1:200 and staining in paraffin-embedded human cervical cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western BlotPositive WB detected in: SH-SY5Y whole cell lysate, U87 whole cell lysate, K562 whole cell lysate, MCF-7 whole cell lysate, HepG2 whole cell lysateAll lanes: INSRR antibody at 1:2000SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 144 kDaObserved band size: 144 kDa

ADDITIONAL INFORMATION

Product Categories/Family for anti-INSRR antibody	Signal Transduction
---	---------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	31657140
NCBI GeneID	3645
NCBI Accession #	NP_055030.1
NCBI GenBank Nucleotide #	NM_014215.2
UniProt Accession #	P14616; O60724; Q5VZS3
Molecular Weight	143,720 Da
NCBI Official Full Name	insulin receptor-related protein
NCBI Official Synonym Full Names	insulin receptor-related receptor
NCBI Official Symbol	INSRR
NCBI Official Synonym Symbols	IRR
NCBI Protein Information	insulin receptor-related protein; IR-related receptor
UniProt Protein Name	Insulin receptor-related protein
UniProt Gene Name	INSRR
UniProt Synonym Gene Names	IRR; IRR
UniProt Entry Name	INSRR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/insrr-antibody-immunohistochemistry-western-blot-elisa/243554>