

Ephrin-A5 (EFNA5) Active Protein | EFNA5 active protein

Recombinant Human Ephrin-A5 (EFNA5) (Active)

CATALOG #
AAA243728


HOST
Mammalian cell

PRODUCT SYNONYMS


Ephrin-A5 (EFNA5); Recombinant Human Ephrin-A5 (EFNA5) (Active); AL-1; EPH-related receptor tyrosine kinase ligand 7; LERK-7; EFNA5 active protein

PRODUCT SPECIFICATIONS

Host	Mammalian cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized from a 0.2 um filtered PBS, 6% Trehalose, pH 7.4.
Sequence Positions	21-203aa; Full Length of Mature Protein
Sequence	QDPGSKAVAD RYAVYWSSN PRFQRGDYHI DVCINDYLDV FCPHYEDSVP EDKTERYVLY MVNFDGYSAC DHTSKGFKRW ECNRPHSPNG PLKFSEKFQL FTPFSLGFEF RPGREYFYIS SAIPDNGRRS CLKLKVFVRP TNSCMKTIGV HDRVFDVNDK VENSLEPADD TVHESAEPSR GEN
Sequence Length	228
Species	Human
Tag	C-terminal FC-tagged
Endotoxin	Less than 1.0 EU/ug as determined by LAL method.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized EPHA3 at 2 ug/ml can bind human EFNA5, the EC50 of human EFNA5 protein is 0.8674-1.119 ng/ml.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Preparation and Storage	Repeated freezing and thawing is not recommended. Store working aliquots at 4 degree C for up to one week.



Bioactivity



SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Related Product Information for EFNA5 active protein	Cell surface GPI-bound ligand for Eph receptors, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Induces compartmentalized signaling within a caveolae-like membrane microdomain when bound to the extracellular domain of its cognate receptor. This signaling event requires the activity of the Fyn tyrosine kinase. Activates the EPHA3 receptor to regulate cell-cell adhesion and cytoskeletal organization. With the receptor EPHA2 may regulate lens fiber cells shape and interactions and be important for lens transparency maintenance. May function actively to stimulate axon fasciculation. The interaction of EFNA5 with EPHA5 also mediates communication between pancreatic islet cells to regulate glucose-stimulated insulin secretion. Cognate/functional ligand for EPHA7, their interaction regulates brain development modulating cell-cell adhesion and repulsion.
Product Categories/Family for EFNA5 active protein	Cancer

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503487
NCBI GeneID	1946
NCBI Accession #	NP_001953.1
NCBI GenBank Nucleotide #	NP_001953.1
UniProt Accession #	P52803
Molecular Weight	50.1 kDa
NCBI Official Full Name	ephrin-A5
NCBI Official Synonym Full Names	ephrin A5
NCBI Official Symbol	EFNA5
NCBI Official Synonym Symbols	AF1; EFL5; RAGS; EPLG7; GLC1M; LERK7
NCBI Protein Information	ephrin-A5
UniProt Protein Name	Ephrin-A5
UniProt Gene Name	EFNA5

UniProt Synonym Gene Names	EPLG7; LERK7; LERK-7
UniProt Entry Name	EFNA5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mammalian-cell-active-protein/ephrin-a5-efna5/243728>