

Rabbit anti-Human CAMK2A Polyclonal Antibody | anti-CAMK2A antibody

CAMK2A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243797	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

CAMK2A; Polyclonal Antibody; CAMK2A Antibody; Calcium/calmodulin-dependent protein kinase type II subunit alpha (CaM kinase II subunit alpha) (CaMK-II subunit alpha) (EC 2.7.11.17); CAMKA KIAA0968; anti-CAMK2A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH7.4
Applicable Applications for anti-CAMK2A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Conjugation	Non-conjugated
Immunogen	Recombinant Human Calcium/calmodulin-dependent protein kinase type II subunit alpha protein (321-478AA)
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image diluted at 1:200 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for anti-CAMK2A antibody	CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity (By similarity). Phosphorylates transcription factor FOXO3 on 'Ser-298' (PubMed:23805378). Activates FOXO3 transcriptional activity (By similarity).
Product Categories/Family for anti-CAMK2A antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	25952118
NCBI GeneID	815
NCBI Accession #	NP_741960.1
NCBI GenBank Nucleotide #	NM_171825.2
UniProt Accession #	Q9UQM7; Q9UL21; Q9Y2H4; Q9Y352
Molecular Weight	54,088 Da
NCBI Official Full Name	calcium/calmodulin-dependent protein kinase type II subunit alpha isoform 2
NCBI Official Synonym Full Names	calcium/calmodulin-dependent protein kinase II alpha
NCBI Official Symbol	CAMK2A
NCBI Official Synonym Symbols	CAMKA
NCBI Protein Information	calcium/calmodulin-dependent protein kinase type II subunit alpha; CaMKIIAlpha; CaMK-II alpha subunit; caMK-II subunit alpha; CaM-kinase II alpha chain; CaM kinase II alpha subunit; caM kinase II subunit alpha; calcium/calmodulin-dependent protein kinase
UniProt Protein Name	Calcium/calmodulin-dependent protein kinase type II subunit alpha
UniProt Gene Name	CAMK2A
UniProt Synonym Gene Names	CAMKA; KIAA0968; CaM kinase II subunit alpha
UniProt Entry Name	KCC2A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/camk2a-antibody-immunofluorescence-immunohistochemistry-elisa/243797>