

ROCK2 recombinant antibody

ROCK2 Antibody

CATALOG # AAA243802	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot, ELISA
-------------------------------	-----------------------	--	--

PRODUCT SYNONYMS

ROCK2; Monoclonal Recombinant Antibody; ROCK2 Antibody; Rho-associated protein kinase 2 (EC 2.7.11.1) (Rho kinase 2) (Rho-associated; coiled-coil-containing protein kinase 2) (Rho-associated; coiled-coil-containing protein kinase II) (ROCK-II) (p164 ROCK-2); KIAA0619; ROCK2 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	10E1
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for ROCK2 recombinant antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human ROCK2
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa Cells at 1:250, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeated by 0.2% TritonX-100, and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Rabbit IgG ?H+L?.



WB (Western Blot)

Western BlotPositive WB detected in: HeLa whole cell lysate, MCF-7 whole cell lysate, L02 whole cell lysate, HepG2 whole cell lysate, Mouse Brain whole cell lysate, Rat Brain whole cell lysateAll lanes: ROCK2 antibody at 1:1000SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 161 kDaObserved band size: 161 kDa

ADDITIONAL INFORMATION

Related Product Information for ROCK2 recombinant antibody	Protein kinase which is a key regulator of actin cytoskeleton and cell polarity. Involved in regulation of smooth muscle contraction, actin cytoskeleton organization, stress fiber and focal adhesion formation, neurite retraction, cell adhesion and motility via phosphorylation of ADD1, BRCA2, CNN1, EZR, DPYSL2, EP300, MSN, MYL9/MLC2, NPM1, RDX, PPP1R12A and VIM. Phosphorylates SORL1 and IRF4. Acts as a negative regulator of VEGF-induced angiogenic endothelial cell activation. Positively regulates the activation of p42/MAPK1-p44/MAPK3 and of p90RSK/RPS6KA1 during myogenic differentiation. Plays an important role in the timely initiation of centrosome duplication. Inhibits keratinocyte terminal differentiation. May regulate closure of the eyelids and ventral body wall through organization of actomyosin bundles. Plays a critical role in the regulation of spine and synaptic properties in the hippocampus. Plays an important role in generating the circadian rhythm of the aortic myofilament Ca(2+) sensitivity and vascular contractility by modulating the myosin light chain phosphorylation.
Product Categories/Family for ROCK2 recombinant antibody	Cancer; Cell biology; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	41872583
NCBI GeneID	9475
NCBI Accession #	NP_004841.2
NCBI GenBank Nucleotide #	NM_004850.3
UniProt Accession #	O75116; Q53QZ0; Q53SJ7; Q9UQN5
Molecular Weight	160,900 Da
NCBI Official Full Name	rho-associated protein kinase 2
NCBI Official Synonym Full Names	Rho-associated, coiled-coil containing protein kinase 2
NCBI Official Symbol	ROCK2
NCBI Official Synonym Symbols	ROCK-II
NCBI Protein Information	rho-associated protein kinase 2; p164 ROCK-2; rho-associated, coiled-coil-containing protein kinase II
UniProt Protein Name	Rho-associated protein kinase 2
UniProt Gene Name	ROCK2
UniProt Synonym Gene Names	KIAA0619; ROCK-II
UniProt Entry Name	ROCK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.