

GATA3 recombinant antibody

GATA3 Antibody

CATALOG # AAA243833	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

GATA3; Monoclonal Recombinant Antibody; GATA3 Antibody; Trans-acting T-cell-specific transcription factor GATA-3 (GATA-binding factor 3); GATA3 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	2A9
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for GATA3 recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human GATA3
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human placenta tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western BlotPositive WB detected in: RAW264.7 whole cell lysateAll lanes: GATA3 antibody
at 1:1000SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 48,
49 kDaObserved band size: 48 kDa

ADDITIONAL INFORMATION

Related Product Information for GATA3 recombinant antibody	Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses.
Product Categories/Family for GATA3 recombinant antibody	Epigenetics and Nuclear Signaling ; Developmental biology ; Immunology ; Stem cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	50541959
NCBI GeneID	2625
NCBI Accession #	NP_001002295.1
NCBI GenBank Nucleotide #	NM_001002295.1
UniProt Accession #	P23771 ; Q5VWG7 ; Q5VWG8 ; Q96J16
Molecular Weight	47,916 Da
NCBI Official Full Name	trans-acting T-cell-specific transcription factor GATA-3 isoform 1
NCBI Official Synonym Full Names	GATA binding protein 3
NCBI Official Symbol	GATA3
NCBI Official Synonym Symbols	HDR; HDRS
NCBI Protein Information	trans-acting T-cell-specific transcription factor GATA-3; GATA-binding factor 3
UniProt Protein Name	Trans-acting T-cell-specific transcription factor GATA-3
UniProt Gene Name	GATA3
UniProt Entry Name	GATA3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.