

BCL2L1 recombinant antibody

BCL2L1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243870	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

BCL2L1; Monoclonal Recombinant Antibody; BCL2L1 Antibody; Bcl-2-like protein 1 (Bcl2-L-1) (Apoptosis regulator Bcl-X); BCL2L BCLX; BCL2L1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	5E9
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for BCL2L1 recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human Bcl-XL
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western Blot Positive WB detected in: K562 whole cell lysate, HeLa whole cell lysate, NIH/3T3 whole cell lysate, Rat Brain whole cell lysate, Mouse Brain whole cell lysate. All lanes: Bcl-XL antibody at 1:1000. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 27, 19, 26 kDa. Observed band size: 30 kDa.

ADDITIONAL INFORMATION

Related Product Information for BCL2L1 recombinant antibody	Potent inhibitor of cell death. Inhibits activation of caspases. Appears to regulate cell death by blocking the voltage-dependent anion channel (VDAC) by binding to it and preventing the release of the caspase activator, CYC1, from the mitochondrial membrane. Also acts as a regulator of G2 checkpoint and progression to cytokinesis during mitosis.
Product Categories/Family for BCL2L1 recombinant antibody	Cancer; Cell biology; Metabolism; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502381
NCBI GeneID	598
NCBI Accession #	NP_001182.1
NCBI GenBank Nucleotide #	NM_001191.2
UniProt Accession #	Q07817; Q5CZ89; Q5TE65; Q92976; E1P5L6
Molecular Weight	26,049 Da
NCBI Official Full Name	bcl-2-like protein 1 isoform 2
NCBI Official Synonym Full Names	BCL2-like 1
NCBI Official Symbol	BCL2L1
NCBI Official Synonym Symbols	BCLX; BCL2L; BCLXL; BCLXS; Bcl-X; bcl-xL; bcl-xS; PPP1R52; BCL-XL/S
NCBI Protein Information	bcl-2-like protein 1; apoptosis regulator Bcl-X; protein phosphatase 1, regulatory subunit 52
UniProt Protein Name	Bcl-2-like protein 1
UniProt Gene Name	BCL2L1
UniProt Synonym Gene Names	BCL2L; BCLX; Bcl2-L-1
UniProt Entry Name	B2CL1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.