

ERBB2 recombinant antibody

ERBB2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243876	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

ERBB2; Monoclonal Recombinant Antibody; ERBB2 Antibody; Receptor tyrosine-protein kinase erbB-2 (EC 2.7.10.1) (Metastatic lymph node gene 19 protein) (MLN 19) (Proto-oncogene Neu) (Proto-oncogene c-ErbB-2) (Tyrosine kinase-type cell surface receptor HER2) (p185erbB2) (CD antigen CD340); HER2 MLN19 NEU NGL; ERBB2 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	5F6
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for ERBB2 recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human ErbB2 (HER2)
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa Cells at 1?50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeated by 0.2% TritonX-100, and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Rabbit IgG ?H+L?.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for ERBB2 recombinant antibody	Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.
Product Categories/Family for ERBB2 recombinant antibody	Cancer; Tags & Cell Markers; Immunology; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	54792098
NCBI GeneID	2064
NCBI Accession #	NP_001005862.1
NCBI GenBank Nucleotide #	NM_001005862.1
UniProt Accession #	P04626; Q14256; Q6LDV1; Q9UMK4; B2RZG3; B4DHN3
Molecular Weight	137,910 Da
NCBI Official Full Name	receptor tyrosine-protein kinase erbB-2 isoform b
NCBI Official Synonym Full Names	v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian)
NCBI Official Symbol	ERBB2
NCBI Official Synonym Symbols	NEU; NGL; HER2; TKR1; CD340; HER-2; MLN 19; HER-2/neu
NCBI Protein Information	receptor tyrosine-protein kinase erbB-2; herstatin; p185erbB2; proto-oncogene Neu; c-erb B2/neu protein; proto-oncogene c-ErbB-2; metastatic lymph node gene 19 protein; tyrosine kinase-type cell surface receptor HER2; neuroblastoma/glioblastoma derived on
UniProt Protein Name	Receptor tyrosine-protein kinase erbB-2
UniProt Gene Name	ERBB2
UniProt Synonym Gene Names	HER2; MLN19; NEU; NGL; MLN 19
UniProt Entry Name	ERBB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/erbb2-recombinant-antibody-clone-5f6-immunofluorescence-immunohistochemistry-elisa/243876>