

DMD recombinant antibody

DMD Antibody

CATALOG # AAA243884	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

DMD; Monoclonal Recombinant Antibody; DMD Antibody; Dystrophin; DMD recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	4B10
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for DMD recombinant antibody	IHC (Immunohistochemistry), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human Dystrophin
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human skeletal muscle tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human heart tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for DMD recombinant antibody	Anchors the extracellular matrix to the cytoskeleton via F-actin. Ligand for dystroglycan. Component of the dystrophin-associated glycoprotein complex which accumulates at the neuromuscular junction (NMJ) and at a variety of synapses in the peripheral and central nervous systems and has a structural function in stabilizing the sarcolemma. Also implicated in signaling events and synaptic transmission.
Product Categories/Family for DMD recombinant antibody	Neuroscience; Signal transduction; Stem cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	313104240
NCBI GeneID	1756
UniProt Accession #	P11532; Q02295; Q14169; Q14170; Q5JYU0; Q6NSJ9; Q7KZ48; Q8N754; Q9UCW3; Q9UCW4; E9PDN1
Molecular Weight	3685
NCBI Official Full Name	Dystrophin
NCBI Official Synonym Full Names	dystrophin
NCBI Official Symbol	DMD
NCBI Official Synonym Symbols	BMD; CMD3B; MRX85; DXS142; DXS164; DXS206; DXS230; DXS239; DXS268; DXS269; DXS270; DXS272
NCBI Protein Information	dystrophin
UniProt Protein Name	Dystrophin
UniProt Gene Name	DMD
UniProt Entry Name	DMD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/dmd-recombinant-antibody-clone-4b10-immunohistochemistry-elisa/243884>