

ADRB2 recombinant antibody

ADRB2 Antibody

CATALOG # AAA243895	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

ADRB2; Monoclonal Recombinant Antibody; ADRB2 Antibody; Beta-2 adrenergic receptor (Beta-2 adrenoreceptor) (Beta-2 adrenoceptor); ADRB2R B2AR; ADRB2 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	4C11
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for ADRB2 recombinant antibody	IHC (Immunohistochemistry), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human beta 2 Adrenergic Receptor
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human prostate cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human endometrial cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for ADRB2 recombinant antibody	Beta-adrenergic receptors mediate the catecholamine-induced activation of adenylate cyclase through the action of G proteins. The beta-2-adrenergic receptor binds epinephrine with an approximately 30-fold greater affinity than it does norepinephrine.
Product Categories/Family for ADRB2 recombinant antibody	Cardiovascular; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	4501969
NCBI GeneID	154
NCBI Accession #	NP_000015.1
NCBI GenBank Nucleotide #	NM_000024.5
UniProt Accession #	P07550; O14823; O14824; O14825; O14826; Q4JG18; Q53GA6; Q6GMT4; Q6P4D8; Q8NEQ9; B0LPE4; B2R7X2
Molecular Weight	46,459 Da
NCBI Official Full Name	beta-2 adrenergic receptor
NCBI Official Synonym Full Names	adrenoceptor beta 2, surface
NCBI Official Symbol	ADRB2
NCBI Official Synonym Symbols	BAR; B2AR; ADRBR; ADRB2R; BETA2AR
NCBI Protein Information	beta-2 adrenergic receptor; beta-2 adrenoceptor; beta-2 adrenoreceptor; catecholamine receptor; adrenergic, beta-2-, receptor, surface
UniProt Protein Name	Beta-2 adrenergic receptor
UniProt Gene Name	ADRB2
UniProt Synonym Gene Names	ADRB2R; B2AR; Beta-2 adrenoceptor
UniProt Entry Name	ADRB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/adrb2-recombinant-antibody-clone-4c11-immunohistochemistry-elisa/243895>