

ILF3 recombinant antibody

ILF3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243900	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

ILF3; Monoclonal Recombinant Antibody; ILF3 Antibody; Interleukin enhancer-binding factor 3 (Double-stranded RNA-binding protein 76) (DRBP76) (M-phase phosphoprotein 4) (MPP4) (Nuclear factor associated with dsRNA) (NFAR) (Nuclear factor of activated T-cells 90 kDa) (NF-AT-90) (Translational control protein 80) (TCP80); DRBF MPHOSPH4 NF90; ILF3 recombinant antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	7H8
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for ILF3 recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human ILF3
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa Cells at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeated by 0.2% TritonX-100, and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Rabbit IgG ?H+L?.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human ovarian cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for ILF3 recombinant antibody	May facilitate double-stranded RNA-regulated gene expression at the level of post-transcription. Can act as a translation inhibitory protein which binds to coding sequences of acid beta-glucosidase (GCase) and other mRNAs and functions at the initiation phase of GCase mRNA translation, probably by inhibiting its binding to polysomes. Can regulate protein arginine N-methyltransferase 1 activity. May regulate transcription of the IL2 gene during T-cell activation. Can promote the formation of stable DNA-dependent protein kinase holoenzyme complexes on DNA. The phosphorylated form at Thr-188 and Thr-315, in concert with EIF2AK2/PKR can inhibit vesicular stomatitis virus (VSV) replication (By similarity).
Product Categories/Family for ILF3 recombinant antibody	Epigenetics and Nuclear Signaling; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	62512150
NCBI GeneID	3609
UniProt Accession #	Q12906; O43409; Q6P1X1; Q86XY7; Q99544; Q99545; Q9BZH4; Q9BZH5; Q9NQ95; Q9NQ96; A8K6F2; G5E9M5
Molecular Weight	894
NCBI Official Full Name	Interleukin enhancer-binding factor 3
NCBI Official Synonym Full Names	interleukin enhancer binding factor 3, 90kDa
NCBI Official Symbol	ILF3
NCBI Official Synonym Symbols	CBTF; DRBF; MMP4; MPP4; NF90; NFAR; NF110; NF90a; NF90b; NFAR2; TCP80; DRBP76; NF110b; NFAR-1; TCP110; MPHOSPH4; NF-AT-90
NCBI Protein Information	interleukin enhancer-binding factor 3; M-phase phosphoprotein 4; translational control protein 80; dsRNA binding protein NFAR-2/MPP4; nuclear factor associated with dsRNA; double-stranded RNA-binding protein, 76 kD; nuclear factor of activated T-cells 90
UniProt Protein Name	Interleukin enhancer-binding factor 3
UniProt Gene Name	ILF3
UniProt Synonym Gene Names	DRBF; MPHOSPH4; NF90; DRBP76; MPP4; NFAR; NF-AT-90; TCP80
UniProt Entry Name	ILF3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.