

SLC2A1 recombinant antibody

SLC2A1 Antibody

CATALOG # AAA243903	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

SLC2A1; Monoclonal Recombinant Antibody; SLC2A1 Antibody; Solute carrier family 2; facilitated glucose transporter member 1 (Glucose transporter type 1; erythrocyte/brain) (GLUT-1) (HepG2 glucose transporter); GLUT1; SLC2A1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	2E1
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for SLC2A1 recombinant antibody	IHC (Immunohistochemistry), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human Glucose Transporter GLUT1
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for SLC2A1 recombinant antibody	Facilitative glucose transporter. This isoform may be responsible for constitutive or basal glucose uptake. Has a very broad substrate specificity; can transport a wide range of aldoses including both pentoses and hexoses.
Product Categories/Family for SLC2A1 recombinant antibody	Cancer; Cardiovascular; Metabolism; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	166795299
NCBI GeneID	6513
NCBI Accession #	NP_006507.2
NCBI GenBank Nucleotide #	NM_006516.2
UniProt Accession #	P11166; O75535; Q147X2; A8K9S6; B2R620; D3DPX0
Molecular Weight	54,084 Da
NCBI Official Full Name	solute carrier family 2, facilitated glucose transporter member 1
NCBI Official Synonym Full Names	solute carrier family 2 (facilitated glucose transporter), member 1
NCBI Official Symbol	SLC2A1
NCBI Official Synonym Symbols	PED; DYT9; GLUT; DYT17; DYT18; EIG12; GLUT1; HTLVR; GLUT1DS
NCBI Protein Information	solute carrier family 2, facilitated glucose transporter member 1; GLUT-1; hepG2 glucose transporter; glucose transporter type 1, erythrocyte/brain; human T-cell leukemia virus (I and II) receptor
UniProt Protein Name	Solute carrier family 2, facilitated glucose transporter member 1
UniProt Gene Name	SLC2A1
UniProt Synonym Gene Names	GLUT1; GLUT-1
UniProt Entry Name	GTR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/slc2a1-recombinant-antibody-clone-2e1-immunohistochemistry-elisa/243903>