

HDAC3 recombinant antibody

HDAC3 Antibody

CATALOG # AAA243972	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

HDAC3; Monoclonal Recombinant Antibody; HDAC3 Antibody; Histone deacetylase 3 (HD3) (EC 3.5.1.98) (RPD3-2) (SMAP45); HDAC3 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	4D4
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for HDAC3 recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human HDAC3
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human endometrial cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western Blot Positive WB detected in: HeLa whole cell lysate, HL60 whole cell lysate, U251 whole cell lysate, K562 whole cell lysate, 293T whole cell lysate All lanes: HDAC3 antibody at 1:2000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 49, 50 kDa Observed band size: 49 kDa

ADDITIONAL INFORMATION

Related Product Information for HDAC3 recombinant antibody	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates. Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys-27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (PubMed:21444723, PubMed:23911289). Contributes, together with XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:25190803).
Product Categories/Family for HDAC3 recombinant antibody	Epigenetics and Nuclear Signaling; Cancer; Cardiovascular; Metabolism; Stem cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	13128862
NCBI GeneID	8841
NCBI Accession #	NP_003874.2
NCBI GenBank Nucleotide #	NM_003883.3
UniProt Accession #	O15379; O43268; Q9UEI5; Q9UEV0; D3DQE1
Molecular Weight	48,848 Da
NCBI Official Full Name	histone deacetylase 3
NCBI Official Synonym Full Names	histone deacetylase 3
NCBI Official Symbol	HDAC3
NCBI Official Synonym Symbols	HD3; RPD3; RPD3-2
NCBI Protein Information	histone deacetylase 3; SMAP45
UniProt Protein Name	Histone deacetylase 3
UniProt Gene Name	HDAC3
UniProt Synonym Gene Names	HD3
UniProt Entry Name	HDAC3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/hdac3-recombinant-antibody-clone-4d4-immunohistochemistry-western-blot-elisa/243972>