

PTGS2 recombinant antibody

PTGS2 Antibody

CATALOG # AAA243983	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunofluorescence, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

PTGS2; Monoclonal Recombinant Antibody; PTGS2 Antibody; Prostaglandin G/H synthase 2 (EC 1.14.99.1) (Cyclooxygenase-2) (COX-2) (PHS II) (Prostaglandin H2 synthase 2) (PGH synthase 2) (PGHS-2) (Prostaglandin-endoperoxide synthase 2); COX2; PTGS2 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	10B2
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for PTGS2 recombinant antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human COX2
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HepG2 Cells at 1:250, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeated by 0.2% TritonX-100, and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Rabbit IgG ?H+L?.



WB (Western Blot)

Western BlotPositive WB detected in: U-87 whole cell lysateAll lanes: COX2 Antibody at 1:1000SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 69 kDaObserved band size: 69 kDa

ADDITIONAL INFORMATION

Related Product Information for PTGS2 recombinant antibody

Converts arachidonate to prostaglandin H2 (PGH2), a committed step in prostanoid synthesis. Constitutively expressed in some tissues in physiological conditions, such as the endothelium, kidney and brain, and in pathological conditions, such as in cancer. PTGS2 is responsible for production of inflammatory prostaglandins. Up-regulation of PTGS2 is also associated with increased cell adhesion, phenotypic changes, resistance to apoptosis and tumor angiogenesis. In cancer cells, PTGS2 is a key step in the production of prostaglandin E2 (PGE2), which plays important roles in modulating motility, proliferation and resistance to apoptosis.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506265
NCBI GeneID	5743
NCBI Accession #	NP_000954.1
NCBI GenBank Nucleotide #	NM_000963.2
UniProt Accession #	P35354; Q16876; A8K802
Molecular Weight	68,996 Da
NCBI Official Full Name	prostaglandin G/H synthase 2
NCBI Official Synonym Full Names	prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase)
NCBI Official Symbol	PTGS2
NCBI Official Synonym Symbols	COX2; COX-2; PHS-2; PGG/HS; PGHS-2; hCox-2; GRIPGHS
NCBI Protein Information	prostaglandin G/H synthase 2; PHS II; PGH synthase 2; cyclooxygenase 2b; prostaglandin H2 synthase 2
UniProt Protein Name	Prostaglandin G/H synthase 2
UniProt Gene Name	PTGS2
UniProt Synonym Gene Names	COX2; COX-2; PGH synthase 2; PGHS-2
UniProt Entry Name	PGH2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/ptgs2-recombinant-antibody-clone-10b2-immunofluorescence-western-blot-elisa/243983>