

CD8A recombinant antibody

CD8A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243987	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

CD8A; Monoclonal Recombinant Antibody; CD8A Antibody; T-cell surface glycoprotein CD8 alpha chain (T-lymphocyte differentiation antigen T8/Leu-2) (CD antigen CD8a); MAL; CD8A recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	2F2
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for CD8A recombinant antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human CD8
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IP (Immunoprecipitation)

Immunoprecipitating CD8A in HepG2 whole cell lysate. Lane 1: Rabbit control IgG instead of in HepG2 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000). Lane 2: 3ug + HepG2 whole cell lysate? 500ug? Lane 3: HepG2 whole cell lysate (10ug)




IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human tonsil tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western Blot Positive WB detected in: MCF-7 whole cell lysate, THP-1 whole cell lysate, PC-3 whole cell lysate, 293 whole cell lysate, HepG2 whole cell lysate, A549 whole cell lysate. All lanes: CD8A antibody at 1:2000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 26, 22, 31 kDa. Observed band size: 31 kDa.

ADDITIONAL INFORMATION

Related Product Information for CD8A recombinant antibody	Identifies cytotoxic/suppressor T-cells that interact with MHC class I bearing targets. CD8 is thought to play a role in the process of T-cell mediated killing. CD8 alpha chains binds to class I MHC molecules alpha-3 domains.
Product Categories/Family for CD8A recombinant antibody	Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	225007536
NCBI GeneID	925
NCBI Accession #	NP_001139345.1
NCBI GenBank Nucleotide #	NM_001145873.1
UniProt Accession #	P01732 ; Q13970 ; Q4ZG17 ; D6W5M8
Molecular Weight	32kDa
NCBI Official Full Name	T-cell surface glycoprotein CD8 alpha chain isoform 1
NCBI Official Synonym Full Names	CD8a molecule
NCBI Official Symbol	CD8A
NCBI Official Synonym Symbols	CD8; MAL; p32; Leu2
NCBI Protein Information	T-cell surface glycoprotein CD8 alpha chain; T8 T-cell antigen; T cell co-receptor; OKT8 T-cell antigen; T-cell antigen Leu2; Leu2 T-lymphocyte antigen; CD8 antigen, alpha polypeptide (p32); T-lymphocyte differentiation antigen T8/Leu-2
UniProt Protein Name	T-cell surface glycoprotein CD8 alpha chain
UniProt Gene Name	CD8A
UniProt Synonym Gene Names	MAL
UniProt Entry Name	CD8A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.