

VDR recombinant antibody

VDR Antibody

CATALOG # AAA243989	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

VDR; Monoclonal Recombinant Antibody; VDR Antibody; Vitamin D3 receptor (VDR) (1;25-dihydroxyvitamin D3 receptor) (Nuclear receptor subfamily 1 group I member 1); NR1I1; VDR recombinant antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	3C7
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for VDR recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human Vitamin D Receptor
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western BlotPositive WB detected in: MCF-7 whole cell lysate, THP-1 whole cell lysate, HL-60 whole cell lysateAll lanes: Vitamin D antibody at 1:1000SecondaryGoat polyclonal to rabbit IgG at 1/50000Predicted band size: 49, 54 kDaObserved band size: 49 kDa

ADDITIONAL INFORMATION

Related Product Information for VDR recombinant antibody	Nuclear hormone receptor. Transcription factor that mediates the action of vitamin D3 by controlling the expression of hormone sensitive genes. Recruited to promoters via its interaction with BAZ1B/WSTF which mediates the interaction with acetylated histones, an essential step for VDR-promoter association. Plays a central role in calcium homeostasis.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507883
NCBI GeneID	7421
NCBI Accession #	NP_000367.1
NCBI GenBank Nucleotide #	NM_000376.2
UniProt Accession #	P11473; Q5PSV3; B2R5Q1
Molecular Weight	48,289 Da
NCBI Official Full Name	vitamin D3 receptor isoform VDRA
NCBI Official Synonym Full Names	vitamin D (1,25- dihydroxyvitamin D3) receptor
NCBI Official Symbol	VDR
NCBI Official Synonym Symbols	NR1I1
NCBI Protein Information	vitamin D3 receptor; 1,25-dihydroxyvitamin D3 receptor; vitamin D nuclear receptor variant 1; nuclear receptor subfamily 1 group I member 1
UniProt Protein Name	Vitamin D3 receptor
UniProt Gene Name	VDR
UniProt Synonym Gene Names	NR1I1; VDR
UniProt Entry Name	VDR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/vdr-recombinant-antibody-clone-3c7-immunohistochemistry-western-blot-elisa/243989>