

WT1 recombinant antibody

WT1 Antibody

CATALOG # AAA243991	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

WT1; Monoclonal Recombinant Antibody; WT1 Antibody; Wilms tumor protein (WT33); WT1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	5F2
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for WT1 recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human Wilms Tumor Protein
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western BlotPositive WB detected in: K562 whole cell lysate, 293T whole cell lysate, A549 whole cell lysate, HEK293 whole cell lysateAll lanes: Wilms Tumor Protein antibody at 1:1000SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 50, 48, 49, 35, 57, 56, 34 kDaObserved band size: 50 kDa

ADDITIONAL INFORMATION

Related Product Information for WT1 recombinant antibody	Transcription factor that plays an important role in cellular development and cell survival (PubMed:7862533). Recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3' (PubMed:7862533, PubMed:17716689, PubMed:25258363). Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors (PubMed:15520190). Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing (PubMed:16934801). Isoform 1 has lower affinity for DNA, and can bind RNA (PubMed:19123921).
Product Categories/Family for WT1 recombinant antibody	Epigenetics and Nuclear Signaling; Cancer; Developmental biology; Tags & Cell Markers

NCBI AND UNIPROT INFORMATION

NCBI GI #	65507817
NCBI GeneID	7490
NCBI Accession #	NP_077742.2
NCBI GenBank Nucleotide #	NM_024424.3
UniProt Accession #	P19544; Q15881; Q16256; Q16575; Q4VXV4; Q4VXV5; Q4VXV6; Q8IYZ5; A8K6S1; B3KSA5
Molecular Weight	49,188 Da
NCBI Official Full Name	Wilms tumor protein isoform B
NCBI Official Synonym Full Names	Wilms tumor 1
NCBI Official Symbol	WT1
NCBI Official Synonym Symbols	GUD; AWT1; WAGR; WT33; NPHS4; WIT-2; EWS-WT1
NCBI Protein Information	Wilms tumor protein; amino-terminal domain of EWS last three zinc fingers of the DNA-binding domain of WT1
UniProt Protein Name	Wilms tumor protein
UniProt Gene Name	WT1
UniProt Entry Name	WT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/wt1-recombinant-antibody-clone-5f2-immunohistochemistry-western-blot-elisa/243991>