

HSPD1 recombinant antibody

HSPD1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243993	Rabbit	Human, Mouse	Immunoprecipitation, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

HSPD1; Monoclonal Recombinant Antibody; HSPD1 Antibody; 60 kDa heat shock protein; mitochondrial (EC 3.6.4.9) (60 kDa chaperonin) (Chaperonin 60) (CPN60) (Heat shock protein 60) (HSP-60) (Hsp60) (HuCHA60) (Mitochondrial matrix protein P1) (P60 lymphocyte protein); HSP60; HSPD1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	3D8
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for HSPD1 recombinant antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human Hsp60
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IP (Immunoprecipitation)

Immunoprecipitating HSPD1 in HepG2 whole cell lysate Lane 1: Rabbit control IgG instead of in HepG2 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000) Lane 2: HepG2 whole cell lysate?500ug? Lane 3: HepG2 whole cell lysate (10ug)




IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human liver tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western Blot Positive WB detected in: Hela whole cell lysate, HepG2 whole cell lysate, 293 whole cell lysate, Jurkat whole cell lysate, MCF-7 whole cell lysate, K562 whole cell lysate, Mouse brain tissue. All lanes: HSPD1 antibody at 1:2000. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 62, 18 kDa. Observed band size: 60 kDa.

ADDITIONAL INFORMATION

Related Product Information for HSPD1 recombinant antibody	Chaperonin implicated in mitochondrial protein import and macromolecular assembly. Together with Hsp10, facilitates the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix (PubMed:1346131, PubMed:11422376). The functional units of these chaperonins consist of heptameric rings of the large subunit Hsp60, which function as a back-to-back double ring. In a cyclic reaction, Hsp60 ring complexes bind one unfolded substrate protein per ring, followed by the binding of ATP and association with 2 heptameric rings of the co-chaperonin Hsp10. This leads to sequestration of the substrate protein in the inner cavity of Hsp60 where, for a certain period of time, it can fold undisturbed by other cell components. Synchronous hydrolysis of ATP in all Hsp60 subunits results in the dissociation of the chaperonin rings and the release of ADP and the folded substrate protein (Probable).
Product Categories/Family for HSPD1 recombinant antibody	Isotype>Loading Controls ; Tags & Cell Markers ; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	31542947
NCBI GeneID	3329
NCBI Accession #	NP_002147.2
NCBI GenBank Nucleotide #	NM_002156.4
UniProt Accession #	P10809 ; Q38L19 ; Q9UCR6 ; B2R5M6
Molecular Weight	61,055 Da
NCBI Official Full Name	60 kDa heat shock protein, mitochondrial
NCBI Official Synonym Full Names	heat shock 60kDa protein 1 (chaperonin)
NCBI Official Symbol	HSPD1
NCBI Official Synonym Symbols	HLD4; CPN60; GROEL; HSP60; HSP65; SPG13; HSP-60; HuCHA60
NCBI Protein Information	60 kDa heat shock protein, mitochondrial; chaperonin 60; 60 kDa chaperonin; heat shock protein 65; P60 lymphocyte protein; mitochondrial matrix protein P1; short heat shock protein 60 Hsp60s1
UniProt Protein Name	60 kDa heat shock protein, mitochondrial
UniProt Gene Name	HSPD1
UniProt Synonym Gene Names	HSP60; CPN60; HSP-60; Hsp60
UniProt Entry Name	CH60_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.